

Public-data File 90-29

**EVIDENCE FOR EPISODIC LATE-HOLOCENE SUBSIDENCE IN ESTUARINE
DEPOSITS NEAR ANCHORAGE, ALASKA: BASIS FOR DETERMINING
RECURRENCE INTERVALS OF MAJOR EARTHQUAKES**

By

R.A. Combellick
Alaska Division of Geological & Geophysical Surveys

November 1990

THIS REPORT HAS NOT BEEN REVIEWED FOR
TECHNICAL CONTENT (EXCEPT AS NOTED IN
TEXT) OR FOR CONFORMITY TO THE
EDITORIAL STANDARDS OF DGGS

794 University Avenue, Suite 200
Fairbanks, Alaska 99709-3645

CONTENTS

	Page
Abstract	1
Introduction	1
Field program.....	6
Equipment and logistics.....	6
Sample collection.....	6
Borehole locations and elevations.....	10
Laboratory analyses	10
Core logging and subsampling.....	10
Radiocarbon dating	11
Plant identification	11
Grain-size analyses.....	11
Results.....	14
Lithology and stratigraphy	14
Radiocarbon ages.....	16
Comparison with previous data.....	20
Discussion	22
Sedimentation and subsidence rates.....	22
Paleoseismicity	23
Summary and conclusions	27
Acknowledgments.....	28
References cited	29
Appendix A - Drilling logs.....	32
Appendix B - Borehole sample descriptions	41
Appendix C - Grain-size analyses	59

FIGURES

Figure 1. Map showing tectonic deformation associated with the great Alaska earthquake of March 27, 1964.....	2
2. Aerial photograph of the Portage area, upper Turnagain Arm.....	4
3. Photograph of pre-1964 vegetation and peat buried as a result of submergence during the great Alaska earthquake.....	5
4. Location map of region near Anchorage, Alaska.....	7
5. Photograph of borehole drilling near Goose Bay.....	8
6. Cross section of hollow-stem auger and continuous-sampling system.....	9
7. Lithologic logs of boreholes drilled in 1988 and 1989 in upper Turnagain and Knik Arms.....	15
8. Photographs of buried peat layers in boreholes TA1 and TA8 at Portage.....	17
9. Plots of age versus depth for six boreholes in upper Turnagain and Knik Arms	19
10. Comparison between radiocarbon ages of samples from boreholes drilled at adjacent locations near Girdwood and Portage in 1985 and 1988.....	21
11. Reported radiocarbon ages of submerged peats and uplifted terraces in the region of the 1964 great Alaska earthquake	26

TABLES

Table	1. Radiocarbon ages of samples from boreholes drilled near Girdwood and Portage in 1985.....	3
	2. Radiocarbon ages of samples from boreholes drilled in vicinity of Turnagain and Knik Arms in 1988 and 1989.....	12
	3. Identification of plant fragments in buried peats and organic-rich layers.....	13

EVIDENCE FOR EPISODIC LATE-HOLOCENE SUBSIDENCE IN ESTUARINE DEPOSITS NEAR ANCHORAGE, ALASKA: BASIS FOR DETERMINING RECURRENCE INTERVALS OF MAJOR EARTHQUAKES

by

R.A. Combellick
Division of Geological and Geophysical Surveys
Alaska Department of Natural Resources
3700 Airport Way
Fairbanks, Alaska 99709

ABSTRACT

Borehole drilling and continuous sampling at nine locations in upper Cook Inlet confirm that a partial record of prehistoric subsidence events, possibly associated with 1964-style major earthquakes, is preserved in estuarine sediments of Turnagain and Knik Arms. Multiple peat layers submerged below present high-tide level and buried by intertidal silt and clay record cycles of supratidal vegetation growth, tectonic subsidence, and rapid burial. Radiocarbon dating of these peat layers provides close maximum ages of coseismic subsidence events responsible for their burial and allows calculation of long-term subsidence rates.

Because of uncertainties in some of the radiocarbon-age data and incomplete correlations with data from other studies in the region, it is not yet possible to construct a reliable regional chronology of events. However, the number of buried peats suggests a minimum of six to eight earthquakes during the past 4,200 radiocarbon yr, which translates to a maximum average recurrence interval of about 525 to 700 yr (590 to 780 yr for about the past 4,700 calendar yr if calculated on the basis of tree-ring calibrated ages). In upper Turnagain Arm, three buried peat layers that date to within 100 yr of each other suggest that recurrence intervals may have been as short as several decades. Radiocarbon ages of shallow submerged peat layers at five locations in both Turnagain and Knik Arms suggest that the most recent major subsidence event prior to the 1964 earthquake was at or before about 500 y. a. The data indicate net subsidence rates of 0.23 to 0.30 cm/yr in upper Turnagain Arm for the past 4,200 radiocarbon yr and 0.13 to 0.17 cm/yr in upper Knik Arm for the past 2,100 radiocarbon yr.

INTRODUCTION

Following the catastrophic great Alaska earthquake of March 27, 1964, geoscientists began to address the question of how soon another damaging earthquake is likely to strike this area of southcentral Alaska, which is home to nearly 60 percent of the state's 500,000 residents. Seismologic monitoring of smaller events has allowed detailed mapping of the Benioff zone and megathrust-interplate boundary and has improved estimates of earthquake-recurrence intervals based on analyses of slip rates and strain release. Considering that the instrumental seismologic record extends only back to about the turn of the century and that the 1964 event is the only known great earthquake in this segment of the Aleutian megathrust during historic time, recurrence estimates of great earthquakes based on these historic data must be regarded as tentative. However, a reliable geologic record of major prehistoric subduction-zone earthquakes in this region for the past several thousand years would complement the historic record and would greatly improve the credibility of recurrence estimates. A chronology of these events would also provide a measure of variability from the average recurrence interval.

The 1964 earthquake caused vertical ground displacements over a 100,000-mi² (260,000-km²) area (fig. 1). This vertical deformation consisted of an asymmetric uplift zone

parallel to the arc and an adjacent landward zone of subsidence (Plafker, 1969). Geologic evidence indicates that similar deformation occurred during the late Holocene, or last 5,000 yr. A total of 131 ft (40 m) of uplift occurred at Middleton Island (fig. 1) during at least five distinct uplift pulses during the last 4,470 yr, based on radiocarbon ages of driftwood and peat on marine uplifted terraces (Plafker and Rubin, 1967). This represents a linear uplift rate of about 0.4 in./yr (1 cm/yr). A terrace is presently forming on Middleton Island due to 10.8 ft (3.3 m) of 1964 uplift. Based on the known ages of uplifted terraces, the recurrence interval for megathrust earthquakes like the 1964 earthquake is 500 to 1,350 yr (Plafker and Rubin, 1967).

Knik and Turnagain Arms, located at the northeastern limit of Cook Inlet in the subsidence zone of the 1964 earthquake (fig. 1), are likely basins in which to find sedimentary evidence of subsidence events associated with major earthquakes. Based on post-1964 survey data, the Knik Arm tidal flats subsided at least 2 ft (0.6 m) during the 1964 Alaska earthquake (Foster and Karlstrom, 1967; Plafker, 1969). Subsidence was due to tectonic lowering of the crust and to compaction of unconsolidated materials underneath the tidal flats. This resulted in extensive tidal flooding and in deposition of intertidal silts over the 1964 soil horizon in at least the southwestern half of the flats.

A similar post-1964 cycle (subsidence, saltwater incursion, vegetation kill, intertidal deposition, and revegetation) has been documented for the upper part of Turnagain Arm southeast of Anchorage (Ovenshine and others, 1976; fig. 2). This cycle for upper Turnagain Arm has restored the sediment surface to its pre-earthquake level (Kachadoorian and Ovenshine, 1984; Bartsch-Winkler and Garrow, 1982). In 1964, this region subsided about 7.9 ft (2.4 m), although as much as 1.8 ft (0.55 m) of rebound occurred following the earthquake (Brown and others, 1977). About 5.2 ft (1.6 m) of intertidal silt has been deposited on the 1964 soil horizon (fig. 3), which is about the amount of deposition to be expected for the net subsidence of the region.

In 1985, geotechnical drilling at two Turnagain Arm sites, one at Girdwood and one at Portage, confirmed the presence of multiple organic layers interbedded with tidal deposits below the 1964 soil horizon. Organic layers are each overlain by 2 to 10 ft (0.6 to 3 m) of fine silty sand, sandy silt, and clayey silt and are interpreted to record episodic prehistoric subsidence in Turnagain Arm similar to that resulting from the 1964 great earthquake (Combellick, 1986). Radiocarbon ages of these organic layers are reported in Table 1. Four peat layers were observed

Table 1. Radiocarbon ages of samples from two boreholes drilled near Girdwood (TA-B1) and Portage (TA-B8) in upper Turnagain Arm in 1985. Calibration to calendar years BP is based on data sets for tree rings of known ages (Linick and others, 1985, 1986; Kromer and others, 1986; Pearson and Stuiver, 1986; Pearson and others, 1986; and Stuiver and Pearson, 1986).

BOREHOLE- DEPTH (ft)	LABORATORY NUMBER	MATERIAL DATED	RADIOCARBON AGE (yr BP)	CALIBRATED AGE (yr BP) 1σ min. (cal. mean ages) 1σ max.
TA-B1-10.0	GX-11346	Sedge peat	1935 ± 150	1710 (1881) 2059
TA-B1-20.0	GX-11347	Wood-sedge peat	2735 ± 145	2749 (2848) 2989
TA-B1-22.0	GX-11348	Wood-sedge peat	2660 ± 100	2739 (2767) 2859
TA-B1-25.5	GX-11349	Wood-sedge peat	3365 ± 155	3459 (3628) 3829
TA-B8-16.0	GX-12199	Sedge peat	1320 ± 340	930 (1277) 1550
TA-B8-21.2	GX-11351	Sedge peat	1915 ± 130	1710 (1870) 2039
TA-B8-25.0	GX-11352	Organic silt	3205 ± 195	3219 (3459) 3683
TA-B8-30.2	GX-11353	Woody peat	2635 ± 145	2549 (2757) 2869
"	GX-11740	(Same sample)	2850 ± 150	2789 (2957) 3209
Weighted average of above two analyses			2739 ± 104	2759 (2849) 2949
TA-B8-35.2	GX-11354	Organic silty clay	3790 ± 175	3295 (4154, 4194, 4222) 4429
TA-B8-41.5	GX-11355	Plant fragments	4240 ± 195	4529 (4841) 50416



Figure 2. Oblique northwest aerial view of the Portage area, Seward Highway, and upper Turnagain Arm. This area subsided about 7.9 ft (2.4 m) during the great Alaska earthquake of 1964, causing extensive tidal flooding. The influx of salt water killed most trees in the central part of the photograph and caused rapid deposition of silt on the submerged flat. Silt deposition and about 1.8 ft (0.55 m) of postseismic rebound have restored this part of the flat to predominantly supratidal conditions (photograph by R.A. Combellick, 1984).



Figure 3. Vegetation and peat layer submerged during the great Alaska earthquake of 1964 and buried beneath approximately 5.2 ft (1.6 m) of post-1964 intertidal silt. Part of the overlying silt layer has been removed by erosion. Exposure is in tidal channel near Girdwood in upper Turnagain Arm (photograph by R. A. Combellick, 1984).

below the 1964 peat to a depth of 26 ft (8 m) at the Girdwood site. Radiocarbon ages of these peats indicate four pre-1964 subsidence events during the past 3.5 Ka. At the Portage site, organic layers to depths up to 41 ft (12.5 m) indicate five pre-1964 subsidence events during the past 4.5 Ka. Because sampling at these sites was not continuous, the possibility of additional buried peat layers means that the number of subsidence events suggested by available data must be considered minimal. Buried peat layers are documented at numerous other sections in the intertidal zone of upper Cook Inlet by Bartsch-Winkler and Schmoll (1987).

The purpose of this project was to drill boreholes with continuous sampling at the same sites drilled at Girdwood and Portage in 1985 and at new sites along Knik Arm to document late-Holocene sedimentation and coseismic subsidence events as completely as possible in two separate basins. This study has confirmed that at least a partial record of prehistoric subsidence events is preserved at the drilled locations in both estuaries. Although a complete record of all observed peats has not yet appeared at any single borehole location, radiocarbon ages of several peats can be correlated among two or more locations and with dated terraces on Middleton Island in the 1964 uplift zone.

FIELD PROGRAM

Equipment and Logistics

Boreholes were drilled at nine locations on vegetated supratidal flats, including sites at Girdwood and Portage along Turnagain Arm in September, 1988, and at Goose Bay and Palmer Hay Flats along Knik Arm in January and February, 1989 (fig. 4). All boreholes were drilled using a truck- or track-mounted CME-75 drill rig operated by the Alaska Department of Transportation and Public Facilities (fig. 5). The 40,000-lb reaction load of this drill rig allowed the sampler to be pushed into the sediment without the use of a hammer, reducing artificial sample compaction and disturbance. Because all Knik Arm boreholes were drilled at off-road sites on wetlands within state game refuges, permit stipulations required drilling from a track vehicle during winter when the ground was frozen and snow covered to minimize surface disturbance.

Sample Collection

Samples were collected using a CME continuous-sample-tube system in conjunction with the hollow-stem auger (fig. 6). The sampler consisted of a 5-ft-long (1.5-m), 3.25-in. (8.3-cm) I.D. split steel barrel connected to the drill rig through the auger with non-rotating rods. Two clear plastic liners, each 2.5 ft (76 cm) long and 2.5 in. (6.4 cm) diam were placed end-to-end in the barrel. The cutting head of the sampler extended a short distance ahead of the lead auger, which rotated around the sampler. Drilling and sampling occurred simultaneously as the sampler was pushed into the sediment 5 ft (1.5 m) at a time. Samples were retrieved by lifting the sampler to the surface through the hollow-stem auger, disassembling the split barrel, and removing the filled sample liners.

As the full sample liners were retrieved from the sampler, the ends were immediately covered by plastic caps and sealed with electrical tape. Sample material caught in the cutting head was retained in sealed plastic bags. A generalized sample log was prepared in the field in conjunction with the drilling log (appendix A), but detailed sample logging was postponed until the samples could be opened in the laboratory.

Sample recovery ranged from 78 to 94 percent (see appendix A). Two circumstances resulted in occasional incomplete sample recovery. First, if the sampler encountered very firm material, such as dense peat, frozen soil, or a large pebble, over less firm material, occasionally the firm material plugged the cutting head and forced the softer material to the outside of the




Figure 4. Location map of region near Anchorage, Alaska, showing locations of boreholes drilled on supratidal flats in 1988 and 1989.




Figure 5. Borehole drilling at site KA4 near Goose Bay in January 1989 (see fig. 4 for location).




Figure 6. Cross section of hollow-stem auger with continuous-sampling system (solid black). Drilling and sampling occur simultaneously in 5-ft (1.5-m) increments as the non-rotating sample barrel pushes ahead of the auger.

sample barrel as the sampler penetrated farther. Second, when sampling loose wet sand or silt with no clay for cohesion, occasionally part of the sample flowed out of the sampler while lifting it to the surface.

Samples were placed upright in crates for transport to Fairbanks and stored in the crates until they were opened in the laboratory.

Borehole Locations and Elevations

Latitudes and longitudes of borehole locations TA1 at Girdwood and TA8 at Portage were determined by surveys referenced to local monuments with theodolite and electronic distance measurement (EDM). Locations of boreholes KA1 through KA6 at Goose Bay and Palmer Hay Flats were determined visually by reference to aerial photographs and 1:63,360-scale topographic maps. Lack of sufficient local horizontal control precluded a straightforward theodolite survey of these locations. However, local vertical control was sufficient to determine ground-surface elevations at all borehole locations using theodolite and EDM. Borehole locations and elevations are included with the sample log in Appendix B.

Relative elevation of the ground surface to mean higher high water (MHHW) at each borehole location was calculated using the surveyed ground elevation relative to mean sea level (MSL) and published statistics from the nearest tide-gage stations (U.S. Department of Commerce, 1988). The level of MHHW relative to the borehole samples must be known in order to show that the subsurface peat layers are presently below the minimum elevation at which they could have formed and to document the minimum amount of subsidence and sedimentation that has occurred since their submergence and burial.

LABORATORY ANALYSES

Core Logging and Subsampling

Each core sample was split in half by cutting the plastic liner on opposite sides with a small high-speed saw and slicing through the sample with a sharp knife or thin wire. Each sample was kept in its semicircular liner halves, which served as trays for handling and storage. After cleaning and smoothing the surface, the sample was described in detail (appendix B) and photographed at two scales with high-resolution black-and-white film. A tape measure was placed by each sample and registered to the top of the sample so that photographs could be keyed directly to sample logs.

Subsamples for radiocarbon dating and granulometric analyses were collected from one side of the split samples. Intact (archive) halves and the remainder of the subsampled halves were covered by plastic wrap and stored in core boxes for possible future use. Exact locations of all subsamples were recorded in the sample log (appendix B).

Peat subsamples for radiocarbon dating were normally 0.1 ft (3 cm) thick and were taken from the top of each peat layer so that radiocarbon ages approximate the ages of the ground surface when burial occurred. If the peat layer was less than 0.1 ft thick, the entire layer was removed from the sample half to ensure sufficient quantity for dating. A total of 43 subsamples were collected from the nine boreholes for possible radiocarbon dating.

Some subsamples collected for radiocarbon dating were of organic silt or disseminated fine organic matter that may represent relict peat layers or may contain reworked or transported organics. These samples were collected primarily to obtain ages for calculation of sedimentation rates in portions of cores that were devoid of peats. Radiocarbon ages obtained from these sam-

ples are regarded with some suspicion because they may be contaminated by older organic material.

One or more subsamples were collected from each texturally distinct stratigraphic unit for granulometric analysis. These subsamples were also normally 0.1 ft (3 cm) thick. Rare exceptions included layers of a distinct texture that were less than 0.1 ft thick and layers of coarser material for which more material was needed to provide an adequate sample for sieving. A total of 55 subsamples were collected for grain-size analysis.

Following completion of logging, photography, and subsampling, each core-sample half was wrapped in clear plastic and stored in a core box for possible future use.

Radiocarbon Dating

Subsamples collected for radiocarbon dating were dried in an oven at about 75°C, weighed, wrapped in aluminum foil, and sealed in plastic bags. Dry subsample weights ranged from 3.6 g for peat to 149 g for silt with disseminated organic matter. Each subsample was examined macroscopically for contaminants, such as roots of younger plants or transported older organics, which were removed.

Of 43 organic subsamples collected from the cores, 38 were selected for submission to Geochron Laboratories Division, Krueger Enterprises, Inc., for radiocarbon dating. Three of the core samples required acceleration mass spectrometry because they contained too little carbon for conventional gas-proportional age determination. Two additional samples, KA7-3.0 and KA7-3.5, collected along the Matanuska River bank at location KA7 (fig. 4) were also submitted. Sample KA7-3.0 consisted of flattened fragments of wood branches lying parallel to bedding at a depth of 3.0 ft (0.9 m). Sample KA7-3.5 was collected from a thin (<0.1 ft or 3 cm) buried peat at 3.5 ft (1.1 m) below the ground surface.

Reported radiocarbon ages (table 2) are based on the Libby half life for ^{14}C (5570 yr) and referenced to AD 1950. A correction for $^{13}\text{C}/^{12}\text{C}$ was applied to adjust for natural isotopic fractionation. Conventional ages were calibrated to tree-ring ages (tables 1 and 2) using a procedure described by Stuiver and Reimer (1986).

Plant Identification

Twenty two of the organic samples were submitted to the University of Alaska Museum for identification of plant remains comprising the submerged peat layers. Plant identifications are presented in Table 3.

Grain-size Analyses

Subsamples for grain-size analyses were dried at about 75°C and weighed. Subsample weights ranged from 50 g to 290 g. Six coarse-grained subsamples were analyzed with conventional 8-in.-diam (20.3-cm) sieves. The remaining 49 fine-grained subsamples were analyzed with a Micromeritics Sedigraph 5000ET rapid sediment analyzer, which uses a scanning x-ray beam to determine the weight-percent distribution of fine particles from their settling velocities in a solution of known density and viscosity.

Two splits of about 1.5 g were obtained from each fine-grained sample for Sedigraph analysis. Any organic material was removed, using hydrogen peroxide if necessary, and the sample was dispersed in a solution of 0.55 percent sodium hexametaphosphate. Part of each sample was injected into the x-ray cell directly from the dispersed suspension and results of the analysis were

Table 2. Radiocarbon ages of samples from boreholes drilled in vicinity of Turnagain and Knik Arms in 1988 and 1989. Calibration to calendar years BP is based on data sets for tree rings of known ages (Linick and others, 1985, 1986; Kromer and others, 1986; Pearson and Stuiver, 1986; Pearson and others, 1986; Stuiver and Pearson, 1986; and Stuiver and others, 1986).

BOREHOLE- DEPTH (ft)	LABORATORY NUMBER	MATERIAL DATED	RADIOCARBON AGE (yr BP)	CALIBRATED AGE (yr BP) 1σ min. (cal. mean ages) 1σ max.
TA1-6.4	GX-15208	Sphagnum-sedge peat	negative (post-1950)	
TA1-8.8	GX-15209	Woody moss-sedge peat	510 ± 130	480 (532) 660
TA1-12.8	GX-15210	Sedge peat	815 ± 115	670 (723) 907
TA1-19.7	GX-15212	Sphagnum-sedge peat	2100 ± 75	1988 (2066, 2098, 2100) 2285
TA1-21.2	GX-15211	Sedge-wood peat	1875 ± 125	1635 (1829) 1950
TA1-22.6	GX-15213	Woody moss-sedge peat	2755 ± 80	2774 (2856) 2950
TA1-25.5	GX-15214	Sphagnum-sedge peat	3455 ± 145	3559 (3703) 3909
TA1-55.3	GX-15215	Organic silt	10,375 ± 310	(too old to calibrate)
TA8-4.9	GX-15216	Sedge blades & roots	330 ± 150	0* (327, 363, 431) 520
TA8-7.0	GX-15217-AMS	Scattered organics	1030 ± 130	790 (943) 1060
TA8-12.7	GX-15218	Sphagnum-sedge peat	885 ± 120	690 (786) 930
TA8-13.4	GX-15404	Sedge peat	1495 ± 165	1280 (1386) 1550
TA8-20.3	GX-15219	Moss-sedge peat	1695 ± 80	1527 (1605) 1709
TA8-21.5	GX-15405	Sedge peat	2630 ± 80	2740 (2756) 2792
TA8-24.8	GX-15220	Woody moss-sedge peat	2675 ± 80	2749 (2775) 2857
TA8-30.3	GX-15221	Moss-sedge peat	2705 ± 85	2754 (2789) 2876
TA8-36.2	GX-15222	Thin peat	3015 ± 140	2989 (3217, 3233, 3243) 3379
TA8-41.5	GX-15223	Thin peat	4150 ± 130	4455 (4648, 4672, 4721, 4757, 4816) 4859
TA8-50.8	GX-15224	Organic silt	10,730 ± 525	(too old to calibrate)
KA1-1.7	GX-15406	Wood-sedge-moss peat	negative (post-1950)	
KA1B-2.7	GX-15225	Wood-sedge-moss peat	470 ± 70	489 (516) 544
KA1-5.85	GX-15226	Sedge peat	955 ± 75	784 (917) 943
KA1-11.75	GX-15227	Sedge peat	2080 ± 130	1890 (2054) 2307
KA1-18.2	GX-15228	Scattered organics	7240 ± 295	7729 (8037) 8369
KA1-28.2	GX-15229-AMS	Organic silt	8850 ± 120	(too old to calibrate)
KA4-0.7	GX-15230	Wood-sedge peat	negative (post-1950)	
KA4-3.3	GX-15231	Wood-sedge peat	515 ± 75	507 (534) 628
KA4-24.2	GX-15232-AMS	Organic silt	4533 ± 84	4996 (5099, 5141, 5166, 5284) 5317
KA4B-3.1	GX-15407	Wood-sedge peat	145 ± 75	0* (0*, 18, 144, 212, 266) 290
KA4B-5.8	GX-15233	Wood-sedge-moss peat	510 ± 70	507 (532) 623
KA4B-12.7	GX-15234	Organic silt	9255 ± 420	(too old to calibrate)
KA5-5.0	GX-15235	Wood-moss-sedge peat	1080 ± 85	930 (976) 1066
KA6-2.3	GX-15236	Silty wood-sedge peat	185 ± 130	0* (0*, 3, 156, 183, 279) 424
KA6-3.0	GX-15237	Sedge peat	560 ± 70	523 (549) 650
KA6-4.35	GX-15238	Sedge peat	930 ± 115	720 (800, 838, 845, 907) 960
KA6-7.55	GX-15239	Organic silt	2810 ± 200	2749 (2893, 2904, 2934) 3219
KA6-7.85	GX-15240	Woody peat	1800 ± 125	1560 (1723) 1880
KA6-25.3	GX-15241	Organic silt	11,400 ± 720	(too old to calibrate)
KA7-3.0	GX-15466	Wood fragments	520 ± 70	510 (536) 628
KA7-3.5	GX-15465	Wood-sedge peat	495 ± 120	472 (525) 641

0* = negative calibrated age BP (post-1950).

Table 3. Identification of plant fragments in buried peats and organic-rich layers (identifications by A.R. Batten and B.M. Murray, University of Alaska Museum, 1989).

<u>SAMPLE-DEPTH (ft)</u>	<u>COMPOSITION</u>
TA1 - 6.4	Moss-sedge peat (Calliergon and Sphagnum).
TA1 - 8.8	Sedge-moss (Sphagnum?) peat (mostly sedge). Some wood up to 3 mm diameter.
TA1 - 12.8	Sedge peat with some moss and wood.
TA1 - 19.7	Moss (Sphagnum?)-sedge peat with twigs up to 5 mm diameter.
TA1 - 21.2	Well-preserved sedge peat with lots of wood (maximum diameter 3 mm).
TA1 - 22.6	Moss (Sphagnum?)-sedge peat with some wood up to 4 mm diameter.
TA1 - 25.5	Sphagnum peat with a few sedges.
TA8 - 12.7	Sphagnum(?) peat with a few sedges and fine (1-2 mm) woody roots or twigs.
TA8 - 20.3	Silty moss-sedge peat, mostly moss (non-Sphagnum?). No twigs apparent.
TA8 - 24.8	Decomposed moss (non-Sphagnum?)-sedge peat with some woody twigs up to 2 mm diameter.
TA8 - 30.3	Moss-sedge peat (mostly moss, Drepanocladus and Calliergon), with a few woody twigs up to 2 mm diameter.
TA8B - 7.0	Sand with sedge & wood fragments.
KA1 - 5.85	Well-decomposed sedge peat.
KA1 - 11.75	Silt with some sedge-like organic fragments. Possibly some wood fragments.
KA1B - 2.7	Well-preserved sedge peat with abundant wood with bark; twigs maximum diameter 4-5 mm.
KA4 - 3.3	Sedge peat with abundant well-preserved wood with bark (maximum diameter about 1 cm).
KA4B - 5.8	Moss-sedge peat (Sphagnum?) with abundant wood (and bark?) fragments. Well-preserved twigs, maximum diameter 5.5 mm.
KA5 - 5.0	Moss-sedge peat (moss component Tomenthypnum and Sphagnum) with several twigs (maximum 3 mm diameter) with bark and leaf scars. Tomenthypnum is often calcareous.
KA6 - 2.3	Silty sedge peat with wood twigs (maximum diameter 3 mm). No moss apparent.
KA6 - 3.0	Sedge peat with some moss (non-Sphagnum). No wood apparent.
KA6 - 4.35	Sedge peat, possibly with some woody roots. No mosses apparent.
KA6 - 7.85	Fine-grained mineral with wood fragments up to 9 mm diameter, 2 cm long.

recorded automatically by the Sedigraph on chart paper. The graph was then scaled in full or half phi (ϕ) units to compute size distributions in weight percent.

The Sedigraph was calibrated and periodically checked for accuracy with reference powders. A third split was analyzed for any sample when results for the first two splits had an error greater than 5 percent in the size distribution. Mean grain size, standard deviation (sorting), skewness

(asymmetry), and kurtosis (peakedness) were calculated from phi-size distributions using standard procedures for moment measures (Friedman and Johnson, 1982) and are reported with percent gravel, sand, silt, and clay in Appendix C.

RESULTS

Lithology and Stratigraphy

Subsurface sediment encountered in the boreholes is predominantly clayey silt to fine sandy silt, similar to material on the modern intertidal flats (see fig. 7 for graphic logs, appendix B for detailed sample descriptions), although layers of coarser sand and gravel are present in the upper parts of the Girdwood and Portage boreholes. Gravel comprising the upper 6 ft (2 m) of borehole TA1 at Girdwood is probably road fill. Fine to medium sand in the upper 10 ft (3 m) of borehole TA8 at Portage displays more distinct layering, coarser grain sizes, and size grading than the remaining fine-grained samples and is interpreted as fluvial overbank deposits from Portage Creek.

The fine sediment making up the remainder of the borehole samples is mostly structureless to faintly laminated clayey silt and silty clay (see appendix C for graphic logs of grain-size statistics). Contorted bedding is rare and is attributed in most cases to disturbance during sampling and handling. In these cases, contorted bedding is restricted to soft sediment adjacent to the void at the top of the cores and extends downward along the core walls. Strongly contorted bedding that cannot be attributed to disturbance appears only in the bottom 3 ft (1 m) of borehole KA6 in upper Knik Arm at depths of 40.5 to 43.5 ft (12.3 to 13.3 m) (appendix B). No peat layers are clearly associated with contorted bedding at this or any other location.

Subsurface peat layers were encountered at all borehole locations. These peats are composed of fresh-water grasses, mosses, and some wood (table 3). In boreholes KA2, KA3, and KA5 on Palmer Hay Flats and in KA4 and KA4B at Goose Bay, the peat layers all occur within 6 ft (2 m) of the surface, immediately below zones that could not be sampled because of the wet, very loose condition of the soil beneath the frozen ground. Therefore, it is not possible to determine from these samples whether these near-surface peats are buried beneath intervening layers of silt or are parts of a single surface layer of peat, although the latter seems most likely.

Deeper subsurface peats interlayered with silt appear in boreholes TA1 and TA8 in Turnagain Arm, KA1 and KA6 on Palmer Hay Flats, and at section KA7 exposed in the bluff at the mouth of the Matanuska River near KA1. In borehole TA1 at Girdwood, four distinct peat layers 0.3 to 1.3 ft thick (0.1 to 0.4 m) are overlain by 0.3 to 6.5 ft (0.1 to 2.0 m) of nearly organic-free silt, minor sand, and some pebbles. Two additional peat layers at 8.5 and 21.1 ft (2.6 and 6.7 m) depth appear below sample breaks and may or may not be separate layers. The deepest peat layer is at 25.5 ft (7.8 m); all are below present MHHW.

In borehole TA8 at Portage, six peat layers 0.1 to 0.8 ft thick (3 to 24 cm) are overlain by 0.2 to 6.7 ft (0.06 to 2.0 m) of organic-free clayey silt and fine sandy silt. Two additional thin peat seams (less than 0.05 ft or 1.5 cm) in otherwise uniform silt appear at 36.2 and 41.5 ft depth (11.0 and 12.7 m) and may also represent former supratidal surfaces. All subsurface peats in this borehole are below present MHHW.

Borehole KA1 on Palmer Hay Flats contains two thin peats overlain by silt below the surface organic layer. These peats occur at 5.8-6.0 ft (1.77-1.83 m) and 11.7-12.0 ft (3.57-3.66 m) below present MHHW. Borehole KA6 contains three well-developed peats interlayered with silt at 3.0-3.2 ft (0.91-0.98 m), 4.3-4.6 ft (1.31-1.40 m), and 7.8-7.9 ft (2.38-2.41 m). An additional 0.05-ft or 1.5-cm-thick organic-rich layer that appears to be a weakly developed peat is present at 7.5-ft (2.3-m) depth. The uppermost peat layer is about at the level of present MHHW.




Figure 7. Lithologic logs of boreholes and sections in vicinity of upper Turnagain and Knik Arms. See Table 2 for laboratory numbers of radiocarbon ages.

A 6-ft-high (1.8-m) bluff section near the mouth of the Matanuska River at the edge of Palmer Hay Flats (location KA7 on fig. 4) is composed entirely of silt with scattered organics and roots near the surface and has a layer of compressed wood fragments at 3.0 ft (0.9 m) and a thin layer of peat at 3.5 ft (1.1 m). These organic layers are slightly above MHHW. The peat layer was initially thought to represent the pre-1964 vegetation, because its depth corresponds approximately to the amount of subsidence documented in this area for the 1964 earthquake. However, radiocarbon dating places both the wood layer and peat at an uncalibrated age of about 530 yr BP, corresponding to basal ages obtained for near-surface peats in several other boreholes.

Stratigraphic relations of several buried peat layers indicate that their development and subsequent burial began with gradual shoaling and sparse vegetation growth, then dense vegetation and peat development, followed by sudden burial. Examples are peat layers at 13 ft (4.0 m) and 25.5 ft (7.8 m) in borehole TA1 and at 30.5 ft (9.3 m) in borehole TA8 (fig. 8). Silt deposits below these peats show a gradual increase in plant and root material in an upward direction, a gradational contact at the base of the peat, and a sharp upper contact between the peat and overlying silt, which is nearly devoid of organics or has only scattered sedge blades or twigs. Scattered organics and a large twig in silt overlying the peat at 13 ft in TA1 (fig. 8A) may be analogous to the buried pre-1964 vegetation preserved in tidal channels near Girdwood. These deposits of Placer River Silt (Ovenshine and others, 1976) encase the branches of small bushes that were growing on the supratidal flat at the time of the earthquake on March 27, 1964. Silt immediately overlying the peat at 25.5 ft contains negligible organics but shows a gradual upward increase in organics within about 1 ft (0.3 m) toward the base of the next higher peat layer at 23 ft depth (7 m)(fig. 8B).

In addition to the peats, layers of organic silt were encountered in the boreholes that may or may not represent buried remnants of former supratidal surfaces. These include zones of scattered plant fragments or dark gray, disseminated fine organic matter embedded in silt (fig. 7). Because of the strong possibility that this organic material was retransported, I assume that these layers are not remnants of peats and use their radiocarbon ages only for calculation of maximum ages and approximate sedimentation rates.

Radiocarbon Ages

Radiocarbon ages of selected peats and organic-rich layers are shown in relation to borehole stratigraphy in Figure 7. All ages are reported in uncalibrated radiocarbon years before present (yr BP), referenced to A.D. 1950. The most significant effect of tree-ring calibrations is to increase the ages of the oldest samples (> 3000 yr) by up to several hundred years (tables 1 and 2). As these differences are probably not significant relative to other sources of error and have no major effect on the conclusions of this report, uncalibrated ages are used in figures to simplify comparisons.

In borehole TA1 at Girdwood, the shallowest subsurface peat layer has a negative radiocarbon age (post-1950) and is assumed to be the surface vegetation that was buried following submergence during the 1964 great Alaska earthquake. Its depth at this location is about the same as the known pre-1964 peat in nearby tidal channels. Dated peat layers below the pre-1964 peat yield radiocarbon ages of 510 ± 130 , 815 ± 115 , $2,100 \pm 75$, $1,875 \pm 125$, $2,755 \pm 80$, and $3,455 \pm 145$ yr BP (table 2; fig. 7). A thin layer of organic silt at 55.3 ft (16.9 m) has a radiocarbon age of $10,375 \pm 310$ yr BP, providing a minimum age for deglaciation comparable to the $10,180 \pm 350$ -yr age obtained for a sample of compressed wood and peat in a gravel pit at Girdwood (Bartsch-Winkler and Schmoll, 1984a).

The 510-yr date at 8.8 ft (2.7 m) in TA1 is for a sample collected at the base of a peat layer that is immediately below a sample break. It is not possible to determine from these samples whether this peat is a separate layer or is continuous with the peat at 6.4-7.4 ft (2.0-2.3 m).




Figure 8. Buried peat layers in boreholes TA1 and TA8, showing gradual upward increase in roots and plant fragments below the peats, sharp upper contacts, and nearly organic-free silt and clay overlying the peats. A, peat layer at 13-ft depth (4.0 m) in borehole TA1 (radiocarbon age 815 ± 115 yr). Note twig at 12.1 ft (arrow). B, peat layer at 25.5-ft depth (7.8 m) in borehole TA1 (radiocarbon age $3,455 \pm 145$ yr). Upper part of photo shows gradational lower portion of peat layer at 23 ft (radiocarbon age $2,755 \pm 80$ yr). C, peat layer at 30.5-ft depth (9.3 m) in borehole TA8 (radiocarbon age $2,705 \pm 85$ yr).

The 1,875-yr peat layer at 21 ft (6.4 m) in TA1 is also immediately below a sample break and is discordant with the ages of peats above and below. A plot of age versus depth (fig. 9A) shows that this sample's age is inconsistent with the long-term sedimentation rate indicated by the other radiocarbon ages. The disturbed top of this core sample and the anomalously young age suggest that the sample may be contaminated by younger organic material. Although a separate peat layer may be present at this depth, the age of sampled material is clearly anomalous. Because recovery was not continuous in this sample interval, it is not clear whether there are two separate peats or one continuous peat between 19.7 and 22.2 ft (6.0-6.8 m).

Radiocarbon ages were obtained for 11 layers in borehole TA8 at Portage, including six peat layers greater than 1 in. (2.5 cm) thick, two thin peat seams, and three layers of organic silt. The peats yield ages of 885 ± 120 , $1,495 \pm 165$, $1,695 \pm 80$, $2,630 \pm 80$, $2,675 \pm 80$, $2,705 \pm 85$, $3,015 \pm 140$, and $4,150 \pm 130$ (table 2; fig. 7). An age discordance occurs between the organic-rich layer at 7.0 ft (2.1 m) and the peat layer at 12.6-13.3 ft (3.8-4.0 m). Although either of these ages could be wrong, I regard the 1,030-yr age of the layer at 7.0 ft with suspicion because its organic content is very low, requiring accelerator mass spectrometry (AMS), and the fine-grained organic matter may have been retransported. Also, the 885-yr age at 12.7 ft (3.9 m) correlates closely with the 815-yr age at about the same depth in borehole TA1.

A radiocarbon age of $10,730 \pm 525$ yr BP for organic silt at 50.8 ft (15.5 m) near the base of borehole TA8 at Portage is comparable to the lowermost age obtained at TA1 at Girdwood and provides a minimum age of deglaciation of the Portage area. Previously, the oldest radiocarbon age obtained in the Portage area was $8,230 \pm 100$ yr BP on wood fragments at the base of a 93-m core about 1 mi (1.6 km) northwest of TA8 (Bartsch-Winkler and others, 1983). The discrepancy between the depths of these samples and their ages casts doubt on the validity of one of the ages; the fact that the shallower sample in TA8 gives the oldest age and results in a sedimentation rate about one seventh that of the remainder of the section may be reason to suspect that the 10,730-yr age is erroneous and may represent transported material. However, the 10,730-yr age correlates closely with the 10,375-yr layer at about the same depth in borehole TA1 at Girdwood (fig. 7), so the question of which date is invalid remains unresolved. Even if the 10,730-yr organics in TA8 were retransported, the date is still a valid minimum age for deglaciation at Portage because the site had to be ice-free at the time the organics were deposited.

In borehole KA1 (Palmer Hay Flats), a thin (< 1 ft or 30 cm) layer of organic silt separates peat at the surface from a buried peat at 1.7-2.8 ft (0.52-0.85 m) (fig. 7, appendix B). A sample at the top of this buried peat layer yielded a negative radiocarbon age (post-1950), and a sample at its base dated at 470 ± 70 radiocarbon yr BP. This latter age correlates closely with ages obtained on layers of wood fragments and peat in nearby tidal-channel section KA7 (fig. 4 for location), which dated at 520 ± 70 and 495 ± 120 radiocarbon yr BP (table 2). Deeper buried peat layers at 5.8-6.0 ft (1.77-1.83 m) and 11.7-12.0 ft (3.57-3.66 m) in KA1 yielded radiocarbon ages of 955 ± 75 and $2,080 \pm 130$ yr BP, respectively. Thin layers of organic silt at 18.2 and 28.2 ft (5.55 and 8.60 m) yielded radiocarbon ages of $7,240 \pm 295$ and $8,850 \pm 120$ yr BP, respectively, but may represent transported material.

No buried peat layers were encountered below the surface peat in boreholes KA2 and KA3 on Palmer Hay Flats (fig. 4; fig. 7), so no radiocarbon ages were obtained. Nearby, in borehole KA5, it was not possible to sample the upper 4.7 ft (1.43 m) of soft, wet peat below the ice. However, a basal peat sample at 5.0 ft (1.52 m) yielded a radiocarbon age of $1,080 \pm 85$ yr BP. This may be the age of the base of the surface peat layer or may be the age of a buried peat that correlates with the peat at 5.8 ft (1.77 m) in KA1.

Three well-developed buried peats were encountered in borehole KA6 on Palmer Hay Flats at 3.0, 4.3, and 7.8 ft (0.91, 1.31, and 2.38 m), yielding ages of 560 ± 70 , 930 ± 115 , and $1,800 \pm 125$ radiocarbon yr BP, respectively (fig. 7). A 185 ± 130 -yr age for organic matter in silt




Figure 9. Plots of radiocarbon age versus depth for six boreholes in the vicinities of upper Turnagain and Knik Arms. Crosses (+) indicate age and bars (-) delimit laboratory standard deviation. Sedimentation rates given in cm per radiocarbon yr. Broken lines denote sedimentation rates spanning periods of questionable radiocarbon age.

at 2.3 ft (0.70 m) at the top of the first core probably represents surface peat mixed with silt through sample disturbance. Layered organic silt at 7.55 ft (2.30 m) yields a radiocarbon age of $2,810 \pm 200$ yr BP, which is discordant with ages above and below (fig. 9F), but correlates closely with ages of well-developed peats in boreholes TA1 and TA8 at Girdwood and Portage. A deeper lamina of organic silt at 25.3 ft (7.71 m) dates at $11,400 \pm 720$ yr BP, which, if valid, provides a new minimum age at this location for retreat of glacial ice after the late-Wisconsin Elmendorf advance (Reger and Updike, 1983).

At borehole KA4 (Goose Bay), a negative radiocarbon age (post-1950) was obtained for peat at the base of the recovered portion of surface organics (fig. 7). A sample break at 1.2-3.0 ft (0.37-0.91 m) makes it difficult to determine whether peat at 3.0-3.35 ft (0.91-1.02 m) is the base of the surface organic layer or is buried under intervening silt. The age of a sample at the base of this layer is 515 ± 75 radiocarbon yr BP. A similar condition exists in the upper 6 ft (1.8 m) of nearby borehole KA4B, where an age of 145 ± 75 radiocarbon yr BP was obtained for a sample at 3.1 ft (0.95 m) midway in the surface(?) peat layer and an age of 510 ± 70 radiocarbon yr BP was obtained for a sample at the base of a peat at 5.8 ft (1.77 m), which may be the base of the surface organic layer or may be buried under intervening silt.

No buried peat layers are present below surface peats in boreholes KA4 and KA4B, although there is a possibility that buried peats are deeper than the drilled depth of about 27 ft (8.2 m). Drilling was suspended at this depth because of -30°F temperatures and the resulting problems with drilling equipment. An AMS radiocarbon age of $4,533 \pm 84$ yr BP was obtained for organic silt at 24.2 ft (7.4 m) in borehole KA4, and a conventional radiocarbon age of $9,255 \pm 420$ yr BP was obtained at 12.7 ft (3.9 m) in borehole KA4B. Buried peat layers have been observed below MHHW in a nearby tidal-channel section at depths of about 12 and 25 ft (3.7 and 7.6 m) and are dated at $1,595 \pm 75$ and $3,270 \pm 90$ radiocarbon yr BP, respectively (Bartsch-Winkler and Schmoll, 1984b).

Comparison with Previous Data

A comparison of radiocarbon ages from boreholes drilled in 1988 at Girdwood and Portage with data obtained at the same locations in 1985 (Combellick, 1986) shows some close agreement and some serious disagreement between the data sets (fig. 10). Some peat layers that appear at the same depth in adjacent boreholes correlate very closely in age, but others disagree by several multiples of their laboratory standard deviations. For example, reported ages of the peat layer at about 25-ft (7.6 m) depth in adjacent Girdwood boreholes differ by only 90 radiocarbon yr, well within the $\pm 1\sigma$ range of both samples. In contrast, the reported $1,935 \pm 150$ yr age of peat at about 10 ft (3 m) in the 1985 borehole is clearly inconsistent with the much younger reported ages (510 ± 130 and 815 ± 115 yr) of peats that appear slightly above and below it in the adjacent 1988 borehole. Similar discrepancies exist between adjacent boreholes at Portage, although peats at 30.3 ft and 41.5 ft (9.2 and 12.7 m) correlate very closely (fig. 10).

Peat at 10-ft (3 m) depth in the 1985 and 1988 Girdwood boreholes is at about the same level (taking into account earthquake subsidence in 1964) as a peat layer dated by Karlstrom (1964) at 700 ± 250 radiocarbon yr BP. This relation supports the validity of the 510- and 815-yr ages in the 1988 borehole and suggests that the 1935-yr age in the 1985 borehole may be erroneous. Bartsch-Winkler and Schmoll (1987) obtained a radiocarbon age of $1,840 \pm 50$ yr BP for a peat sample collected by auger at a depth of about 22 ft (6.7 m) below the surface in the middle intertidal zone at Girdwood, which may correlate with one of the layers in the 20-23-ft (6.1-7.0 m) depth range in Girdwood boreholes TA-B1 and TA1 but does not clearly resolve the age discrepancies at that depth (fig. 10).

Until additional data are obtained at the Girdwood and Portage sites, the remaining age discrepancies may remain unresolved. Possible reasons for the discrepancies include (1) strati




Figure 10. Comparison between radiocarbon ages of samples from boreholes drilled at adjacent locations near Girdwood and Portage in 1985 and 1988. See Tables 1 and 2 for laboratory numbers.

graphic discontinuities between adjacent boreholes; (2) labeling errors; (3) contamination by external organic matter or bacterial activity; (4) handling errors in the laboratory; or (5) analytical errors arising from faulty equipment, poor technique, or calibration problems.

Tidal or fluvial channel migration disrupts stratigraphic continuity and is evident in numerous other boreholes in the area. However, channel deposits observed in other boreholes are much coarser than the host tidal deposits and are devoid of peat layers. Laboratory errors seem highly unlikely considering that both sets of samples were analyzed at the same reputable laboratory.

Contamination by bacterial activity is a possible source of error in 1988 samples, considering that several months elapsed before subsamples could be submitted for radiocarbon dating. Bacterial uptake of recent carbon dioxide is known to be responsible for anomalously young radiocarbon ages in marine cores stored unfrozen for several years in unsealed core boxes. Geyh and others (1974) showed that the effects of bacterial contamination become more pronounced with age of the sediment, producing highly erratic results on samples of Pleistocene age (more than 10,000 yr old). In all cases, contamination results in a radiocarbon age that is younger than expected. Severely contaminated samples of Holocene age can be in error by up to 2,500 radiocarbon yr.

Unlike the results documented by Geyh and others (1974), results of this study at Girdwood and Portage show the greatest discrepancies in the youngest samples (< 3,000 radiocarbon yr BP). Additionally, although most of the apparent discrepancies in 1988 samples are due to radiocarbon ages that are younger than expected, some are older. Other radiocarbon ages, particularly for older samples, show close agreement with the 1985 results.

Regardless of sources of error, these discrepancies may weaken the significance of exact ages of peat layers for constructing a history of events until a sufficiently large regional data base exists. To avoid confusion and for the sake of consistency, the remaining discussion on sedimentation rates, subsidence, and earthquake implications is limited to data obtained from 1988 and 1989 boreholes. All radiocarbon dating of these samples was performed at the same laboratory at about the same time, which should eliminate any significant effect of differences in laboratory methods.

DISCUSSION

Sedimentation and Subsidence Rates

Average sedimentation rates were computed from the 1988/89 data for intervals between radiocarbon-dated layers and for each borehole section between the oldest dated layer and the surface (fig. 9). If the model for restoration of Portage Flats after the 1964 earthquake (Kachadoorian and Ovenshine, 1984) applies to previous subsidence events, actual sedimentation rates are not constant between events but are very rapid immediately after an event, tapering to very slow or zero for most of the remaining time before the next event. Consequently, reported sedimentation rates between dated layers represent the average rates over periods during which actual rates may have varied considerably.

At location TA1 (Girdwood), the data suggest an overall rate of 0.16 cm/yr for about the past 10,400 radiocarbon yr. With exception of the anomaly caused by the outlier at 21.2 ft (6.5 m)(fig. 9A), the net rate remained relatively constant between 0.13 and 0.40 cm/yr prior to 1964.

At location TA8 (Portage), the data suggest a similar average sedimentation rate of 0.14 cm/yr for the past 10,730 radiocarbon yr. However, this rate is based on a questionable age for the oldest dated sample and is considerably slower than the rate determined by Bartsch-Win-

kler and others (1983) at a nearby borehole. If the rate is computed from the oldest, presumably reliably dated peat at this location ($4,150 \pm 130$ yr BP at 41.5 ft or 12.7 m), the long-term sedimentation rate would be 0.30 cm/yr. This compares favorably to the 0.25-cm rate determined by Bartsch-Winkler and others (1983) for the past 5,740 radiocarbon yr.

Aside from the possible error in the 10,730-yr age, the remaining radiocarbon ages in borehole TA8 suggest that sedimentation rates at this site were more variable than at TA1 (fig. 9B). In particular, the narrow age span of three peat layers between 30.3 and 21.5 ft (9.2 and 6.5 m) suggests sedimentation rates of 0.58 to 5.59 cm/yr, up to 40 times the long-term rate, between $3,015 \pm 140$ and $2,630 \pm 80$ radiocarbon yr BP. Higher sedimentation rates during this period may have resulted from increased glacial activity during the Neoglacial maximum, which was under way in southern Alaska by about 3,000 y.a. and began to recede by about 2,000 y.a. (Calkin, 1988). Close proximity of the Portage site to several large glaciers could explain why sedimentation rates were much higher there than at Girdwood during this period, even though both sites remained dominantly intertidal (based on lack of significant differences in texture between boreholes or above and below the peat layers in this time span). Considering discrepancies in radiocarbon ages between the 1985 and 1988 boreholes at Portage, however, these variations in sedimentation rate may also be an artifact of dating errors.

Long-term sedimentation rates on Palmer Hay Flats in upper Knik Arm, represented by boreholes KA1 and KA6 (figs. 9C and F), appear to be about half of the rate computed for Portage and Girdwood in upper Turnagain Arm. At Goose Bay, the long-term sedimentation rate indicated by radiocarbon dates at KA4 is similar to the Turnagain Arm rate (0.16 cm/yr), but the rate at borehole KA4B is significantly lower (0.04 cm/yr). Considering the close proximity of the Goose Bay boreholes on the same supratidal surface (fig. 4), the sedimentation rate should be nearly the same. Most likely, either the 4,533-yr date in KA4 or the 9,255-yr date in KA4B is in error.

Assuming tidal sedimentation and postseismic rebound restore tidal flats to subaerial conditions between deposition of peat layers, the average sedimentation rate between peat layers is also a measure of the net rate of subsidence. Accordingly, sedimentation rates calculated from dating of the oldest peat layers suggest a net rate of subsidence of 0.23 to 0.30 cm/yr in upper Turnagain Arm for the past 3,400 to 4,200 radiocarbon yr, and 0.13 to 0.17 cm/yr in upper Knik Arm for the past 1,800 to 2,100 radiocarbon yr. The lower subsidence rate for upper Knik Arm appears consistent with measurements following the 1964 earthquake, which showed that Palmer Hay Flats subsided about 0.6 m (Plafker, 1969), compared to about 1.5 m of net subsidence in upper Turnagain Arm, taking into account postseismic uplift (Brown and others, 1977).

Paleoseismicity

Stratigraphic evidence of gradual shoaling, vegetation growth, peat development, and rapid submergence and burial below present high-tide level supports the hypothesis that each buried peat layer represents a vegetated supratidal surface that subsided during a major earthquake. Burial occurred rapidly as new sediment was deposited on the submerged flat, probably similar to sedimentation in upper Turnagain Arm that restored the flat to subaerial conditions within 17 yr following the 1964 earthquake (Kachadoorian and Ovenshine, 1984).

It may be argued that the observed stratigraphy is also consistent with burial of the peats by river-flood deposits, followed by submergence. However, with the exception of the peat layer at 19.7 ft (6.0 m) in TA1, which is overlain by pebbly coarse sand, no significant textural differences are evident in deposits overlying buried peats to indicate a change from tidal to fluvial sedimentation.

Gradual submergence by aseismic subsidence can probably be ruled out as a model for formation and burial of peat layers because sufficiently long stillstands would be necessary for peat formation. Also, gradual submergence would subject the surface to erosion and reworking by storm waves and tidal currents, making preservation of peats less likely. Dragging of large ice masses by tidal currents during winter in Turnagain and Knik Arms would be particularly destructive. Temporary eustatic sea-level rises followed by rapid or gradual subsidence could result in preservation of peats below present sea level, but sea-level rise would have to be sufficiently rapid to submerge and bury the peat so that it is not removed by erosion. Global relative sea level has been relatively constant during the past 5,000 yr and probably has not been higher than at present (Clark and others, 1978). Karlstrom (1965) inferred a high sea-level stand between 5,000 and 6,000 y.a., based on radiocarbon dating of coastal bogs and represented by 'elevated' supratidal flats at the mouths of the Kenai, Kasilof, Susitna and Little Susitna Rivers, at Palmer Hay Flats, and at Goose Bay. If this were true, buried peat layers beneath these surfaces should be greater than 5,000 yr old. However, radiocarbon ages of basal surface peat and shallow buried peat layers in this study (fig. 7) show that the surfaces at Palmer Hay Flats and Goose Bay are less than about 500 yr old. The age of a peat layer buried under clayey silt below the supratidal surface near the mouth of the Little Susitna River shows that the surface there is less than 355 ± 75 radiocarbon yr old (lab number I-11,867, R.D. Reger, unpublished data). These surfaces probably are not elevated but represent the high-intertidal depositional surface equivalent to the re-established surface in upper Turnagain Arm (Kachadoorian and Ovenshine, 1984). These surfaces are flooded during extreme high tides and support growth of primarily sedge grass and other vegetation that can withstand periodic flooding by sea water. I know of no other evidence for sea-level fluctuations during the late Holocene in this region apart from relative effects of tectonic subsidence that were sufficient to account for submergence and burial of peats to the extent observed in upper Turnagain and Knik Arms.

Radiocarbon ages of organic material at the tops of submerged peat layers provide maximum ages for subsidence events. If the vegetation was buried by fluvial-flood deposition before submergence, an unknown period of time passed between vegetation growth and submergence; if burial was due to rapid marine deposition after submergence, the radiocarbon age provides a closely limiting maximum age for the event. The lack of textural changes above the peats suggests that burial was by renewed intertidal deposition following submergence rather than by fluvial flooding. In either case, assuming the hypothesis relating submerged peat layers to major earthquakes is correct, then the peats provide a minimum count of earthquakes that have occurred following deposition of the oldest peat. As previously discussed, apparent age discrepancies make attempts at assigning ages to all prehistoric earthquakes risky at this time.

Regardless of exact timing, the number of peats in a borehole can be considered a minimum count of subsidence events during the period in which they were deposited because additional events may have occurred for which no peat layer was preserved. Tidal deposition may not have restored the flat to subaerial conditions prior to every earthquake. As Bartsch-Winkler and Schmoll (1987) have demonstrated, vegetated and nonvegetated zones of the tidal area migrate seaward and landward over time, so preservation of peat layers representing all subsidence events at any one location is highly unlikely.

The largest number of buried peat layers was encountered in borehole TA8 at Portage. Of eight layers present, six are well-developed peats > 1 in. (2.5 cm) thick and two are thin (<0.5 in. or 1.3 cm) peat seams that arguably do not represent former supratidal surfaces. Therefore, the Portage site shows evidence of at least six to eight subsidence events during about the last 4,200 radiocarbon yr (about 4,700 calendar yr)(table 2) prior to 1964. This translates to a maximum average recurrence interval of 525 to 700 yr (very roughly 590 to 780 yr if calculated on the basis of tree-ring calibrated ages from table 2).

Assuming the radiocarbon ages of the peats at TA8 are valid and the interpretation that they represent subsidence during major earthquakes is correct, the data suggest that there were three major (great?) earthquakes within a period of less than 100 yr between $2,705 \pm 85$ and $2,630 \pm 80$ radiocarbon yr ago. If calibrated ages are used, the time span is only 33 yr plus laboratory standard deviations (table 2). Ignoring possible effects of postearthquake compaction, net subsidence during these events was 5.5 ft (1.7 m), 3.3 ft (1.0 m), and 1.2 ft (0.4 m), respectively; subsidence during the first two events was comparable to subsidence during the 1964 event. The occurrence of three large earthquakes within 100 yr appears inconsistent with the long-term average recurrence interval and contradictory to seismic-gap theory, which states that a region along an active plate boundary that has not produced a large earthquake for a long period of time is more likely to be the site of a future large earthquake than adjacent regions that have ruptured more recently (McCann and others, 1979). The theory implies that after a gap-filling earthquake, sufficient time must elapse before stresses can reaccumulate for release in another large earthquake. The seismic-gap theory was used to successfully forecast many large earthquakes, most recently the October 17, 1989 Loma Prieta, California, earthquake (Ward and Page, 1989). However, there are cases in which multiple large earthquakes have occurred in the same rupture zone within an unusually short period. For example, the May 7, 1986 great earthquake ($M_w = 8.0$) in the Andreanof Islands, Alaska, occurred in approximately the same location as a great earthquake on March 9, 1957 ($M_w = 8.6$). The aftershock zone of the 1986 event overlapped a significant portion of the 1957 rupture zone (Ekström and Engdahl, 1989). The time between these events was only 29 yr, even though the recurrence period had been projected at 249 yr on the basis of a time-predictable probabilistic model (Jacob, 1984). Submergence and burial of three well-developed peats with radiocarbon ages spanning less than 100 yr at Portage may be evidence of similar, unusually brief recurrence times for multiple large earthquakes in the 1964 rupture zone about 2,700 radiocarbon yr ago.

It is not possible at this time to relate apparent amounts of subsidence between peat layers to earthquake magnitudes with any degree of confidence. Only if reliable regional data were available to document the areal distribution of deformations associated with individual events could any inferences be made about relative magnitude. For example, differences in depth between peat layers at Portage (borehole TA8) cannot be used to infer differences in earthquake magnitude because the position of this borehole relative to the zero isobase of deformation is not known for each event. During the 1964 event, vertical displacements varied considerably over relatively short distances (Plafker, 1969). Because the position of the zero isobase and the distribution of vertical deformation undoubtedly is not the same in every large earthquake along this rupture zone, the amount of subsidence at Portage could vary substantially during successive earthquakes of similar magnitude.

Perhaps all that can be said about magnitude of prehistoric earthquakes based on available data for this region is that a large-magnitude event is necessary to produce subsidence or uplift that is likely to be preserved in the geologic record. The relation of length of deformation zone to earthquake magnitude for historic subduction-zone events along the Pacific margin suggests a threshold magnitude of about 7, below which measurable deformation does not occur (West and McCrumb, 1988). A higher threshold is probably necessary for deformation to be preserved as uplifted marine terraces or submerged supratidal peat layers.

Data presently available from this and other studies of coseismic uplift and subsidence in the rupture zone of the 1964 great Alaskan earthquake provide strong evidence that several major events occurred during the past 5,000 yr. However, insufficient regional data are available to construct a credible, complete timetable of events. Comparison of data from this study with data from some other studies in the region shows little obvious correlation of radiocarbon ages that are presumably associated with coseismic uplift or subsidence (fig. 11). As discussed above, data from this study in upper Turnagain Arm, from both 1985 and 1988 boreholes, strongly suggest at least one and possibly three events between 2,600 and 2,900 radiocarbon yr BP. However, radio




Figure 11. Reported radiocarbon ages of submerged peats and uplifted terraces possibly associated with major prehistoric earthquakes in the rupture zone of the 1964 great Alaskan earthquake. Bars represent one laboratory standard deviation about the mean radiocarbon age. Plotted data from this study do not include ages of organic silt layers, peat layers with anomalous radiocarbon ages, or peat layers that are below sample breaks where the overlying stratigraphy was not observed.

carbon ages of these peats do not correlate with reported ages from other areas, except for a discordant age of a poorly developed peat in borehole KA6 in upper Knik Arm ($2,810 \pm 200$ yr; not plotted on fig. 11). Either there are major errors in radiocarbon ages in this or other studies, or burial events were restricted to Turnagain Arm and not associated with large interplate earthquakes, or evidence of the earthquakes has not been discovered in the other areas.

Comparison of data from this study at Portage with data at Middleton Island (Pflafer and Rubin, 1978) suggests possible correlation of events at about 1,300-1,500, 3,800, and 4,100-4,400 radiocarbon yr (fig. 11). There also may be regional evidence of events at about 700-1,000, 1,700-2,100, and 3,100-3,400 radiocarbon yr. Close correlation between the ages of a buried peat layer at Pt. Woronzof (Bartsch-Winkler and Schmoll, 1984b) and an uplifted terrace at Middleton Island suggest another possible event at about 2,300-2,400 radiocarbon yr BP. Considering discrepancies in radiocarbon ages that have been discussed previously, these correlations are highly speculative.

Although radiocarbon ages were obtained for some near-surface peats where sample recovery was incomplete, their meaning with regard to recent subsidence history and earthquake occurrence is obscure because their stratigraphic context is unclear at all locations except KA6 and KA7 on Palmer Hay Flats. Shallow peat layers dating at about 500 radiocarbon yr BP at locations TA1, KA1, KA4, and KA4B all appear below sample breaks that could contain either continuous peat or interbeds of tidal sediments. These breaks in sample continuity occurred where the coring device did not retain soft, wet peat and sediment in the shallow subsurface beneath the frozen ground. If peat is continuous between the 500-yr layer and the surface (or between the 500-yr layer and the pre-1964 peat at TA1), 500 yr is a basal age that represents the onset of vegetation on the pre-1964 supratidal flat and is a minimum age for the most recent major pre-1964 earthquake. If the peat is buried beneath tidal silt, 500 yr (calibrated age about 530 calendar yr)(table 2) is a close maximum age for that event.

At KA6, a peat layer with a radiocarbon age of 560 ± 70 yr is overlain by 3 ft (0.9 m) of silty clay and organic silt to the surface of the supratidal flat. At KA7, a peat layer with a radiocarbon age of 495 ± 120 yr is overlain by 3.5 ft (1.1 m) of silt. At both locations, the 500-yr peat layer is above present MHHW, but still below the level of most high spring tides. Although burial may have resulted from regional subsidence, the possibility of burial during a major flood of either Matanuska or Knik Rivers must also be considered. Glacial-outburst flooding occurred almost annually in Knik River between 1918 and 1966 and could be a mechanism for sedimentation on Palmer Hay Flats during a major flood. However, the very fine-grained texture of the deposits overlying the peat is inconsistent with major fluvial-flood deposition except possibly in very quiet back-water conditions. The fact that the age of these peat layers correlate closely with submerged peats at four other locations in Turnagain and Knik Arms strongly suggests a regional subsidence event that was probably associated with the youngest pre-1964 major earthquake, at or before about 500 y.a.

SUMMARY AND CONCLUSIONS

This study confirms that at least a partial record of prehistoric subsidence events, possibly associated with 1964-style major earthquakes, is preserved in estuarine sediments of Turnagain and Knik Arms in upper Cook Inlet. Multiple peat layers submerged below present high-tide levels and buried beneath intertidal silt and clay record cycles of supratidal vegetation growth, tectonic subsidence, and rapid burial during the period of sea-level stability over the past 5,000 yr. Radiocarbon dating of organic material at the tops of these peat layers provides maximum ages for subsidence events responsible for their burial and allows calculation of long-term subsidence rates. Although uncertainties in some radiocarbon ages preclude accurate dating of all events, the buried peats provide a minimum count of earthquakes from which a maximum average recurrence interval can be calculated.

Sedimentation rates computed from radiocarbon ages of the buried peats provide an estimate of the net rate of subsidence, assuming that the peats developed at approximately the level of the modern supratidal surface. The data indicate net subsidence rates of 0.23 to 0.30 cm/yr in upper Turnagain Arm for the past 4,200 radiocarbon yr and 0.13 to 0.17 cm/yr in upper Knik Arm for the past 2,100 radiocarbon yr.

Subsurface stratigraphy at Portage, where the largest number of buried peat layers was encountered in boreholes, shows evidence of six to eight subsidence events during about the past 4,200 radiocarbon yr (4,700 calendar yr) prior to the 1964 great Alaska earthquake. This translates to a maximum average recurrence interval of 525 to 700 radiocarbon yr (about 590 to 780 calendar yr if calculated on the basis of tree-ring calibrated ages). Radiocarbon ages of shallow submerged peat layers in both Turnagain and Knik Arms indicate that the most recent major subsidence event prior to the 1964 earthquake was at or before about 500 y.a. These conclusions are reasonably consistent with data from the Copper River Delta and Prince William Sound regions (Plafker, 1990) that suggest an average recurrence interval of 650-850 yr since about 3,000 yr BP, with about 850 yr since the youngest major pre-1964 earthquake.

Possible evidence at Portage of three earthquakes within 100 yr between $2,705 \pm 85$ and $2,630 \pm 80$ radiocarbon yr BP suggests that although average recurrence times are on the order of centuries, actual recurrence times may have been as short as several decades. However, because of uncertainties in some of the radiocarbon ages, this conclusion is tentative until more regional data on events during this period are available.

Comparison with other radiocarbon ages of features interpreted to represent coseismic subsidence or uplift do not yet provide a clear chronology of pre-1964 events in the region affected by tectonic deformation during the 1964 earthquake. Although some ages roughly correlate, many do not. This poor correlation of regional data may relate to inaccuracies in radiocarbon dating, local effects not associated with widespread tectonic changes during major earthquakes, or a combination of these factors.

ACKNOWLEDGMENTS

Partial funding for this study was provided by the U.S. Geological Survey, Department of the Interior, under National Earthquake Hazards Reduction Program award 14-08-0001-G1582. The Alaska Department of Transportation and Public Facilities (DOTPF) provided and operated the drill rigs used for all borehole drilling. I especially thank Dan Pavay of DOTPF, who arranged for equipment and organized field logistics, and drillers Paul Mitchell and Les Lee for their expert work under adverse winter conditions. Geochron Laboratories Division, Krueger Enterprises, Inc., performed all radiocarbon-age determinations. I thank Richard Reger for his participation in field work and data interpretation, Bret Davidson for assistance in the laboratory, and Roger Alley for surveying borehole elevations. Thanks also to Alan Batten and Barbara Murray of the University of Alaska Museum for identifying plant remains in the peat samples. Richard Reger thoroughly reviewed the manuscript and provided many valuable suggestions for improvement.

The contents of this report were developed under a grant from the U.S. Geological Survey, Department of the Interior. However, those contents do not necessarily represent the policy of that agency, and you should not assume endorsement by the federal government.

REFERENCES CITED

- Bartsch-Winkler, S.R., and Garrow, H.C., 1982, Depositional system approaching maturity at Portage Flats, in Coonrad, W.L., ed., *The U.S. Geological Survey in Alaska: Accomplishments during 1980*: U.S. Geological Survey Circular 844, p. 115-117.
- Bartsch-Winkler, Susan, and Schmoll, H.R., 1984a, Guide to late Pleistocene and Holocene deposits of Turnagain Arm: Anchorage, Alaska Geological Society, 70 p., 1:63,360, 2 sheets.
- _____, 1984b, Bedding types in Holocene tidal channel sequences, Knik Arm, upper Cook Inlet, Alaska: *Journal of Sedimentary Petrology*, v. 54, no. 4, p. 1239-1250.
- _____, 1987, Earthquake-caused sedimentary couplets in the upper Cook Inlet region, in Hamilton, T.D., and Galloway, J.P., eds., *Geologic Studies in Alaska by the U.S. Geological Survey during 1986*: U.S. Geological Survey Circular 998, p. 92-95.
- Bartsch-Winkler, Susan, Ovenshine, A.T., and Kachadoorian, Reuben, 1983, Holocene history of the estuarine area surrounding Portage, Alaska as recorded in a 93 m core: *Canadian Journal of Earth Sciences*, v. 20, no. 5, p. 802-820.
- Brown, L.D., Reilinger, R.E., Holdahl, S.R., and Balazs, E.I., 1977, Postseismic crustal uplift near Anchorage, Alaska: *Journal of Geophysical Research*, v. 82, no. 23, p. 3369-3378.
- Calkin, P.E., 1988, Holocene glaciation of Alaska (and adjoining Yukon Territory, Canada): *Quaternary Science Reviews*, v. 7, no. 2, p. 159-184.
- Clark, J.A., Farrell, W.E., and Peltier, W.R., 1978, Global changes in postglacial sea level: A numerical calculation: *Quaternary Research*, v. 9, p. 265-287.
- Combellick, R.A., 1986, Chronology of late-Holocene earthquakes in southcentral Alaska: Evidence from buried organic soils in upper Turnagain Arm: *Geological Society of America Abstracts with Programs*, v. 18, no. 6, p. 569.
- Ekström, Göran, and Engdahl, E.R., 1989, Earthquake source parameters and stress distribution in the Adak Island region of the central Aleutian Islands, Alaska: *Journal of Geophysical Research*, v. 94, no. B11, p. 15499-15519.
- Foster, H.L., and Karlstrom, T.N.V., 1967, Ground breakage and associated effects in the Cook Inlet area, Alaska, resulting from the March 27, 1964, earthquake: U.S. Geological Survey Professional Paper 543-F, 28 p.
- Friedman, G.M., and Johnson, K.G., 1982, *Exercises in sedimentology*: New York, John Wiley & Sons, 208 p.
- Geyh, M.A., Krumbein, W.E., and Kudrass, H.R., 1974, Unreliable radiocarbon dating of long-stored deep-sea sediments due to bacterial activity: *Marine Geology*, v. 17, p. M45-M50.
- Jacob, K.H., 1984, Estimates of long-term probabilities for future great earthquakes in the Aleutians: *Geophysical Research Letters*, v. 11, no. 4, p. 295-298.
- Kachadoorian, Reuben, and Ovenshine, A.T., 1984, Natural restoration from the effects of the 1964 earthquake at Portage, southern Alaska, in Coonrad, W.L., and Elliott, R.L., eds., *The United States Geological Survey in Alaska: Accomplishments during 1981*: U.S. Geological Survey Circular 868, p. 109-110.

- Karlstrom, T.N.V., 1964, Quaternary geology of the Kenai lowland and glacial history of the Cook Inlet region, Alaska: U.S. Geological Survey Professional Paper 443, 69 p., 7 sheets.
- _____ 1965, Resumé of the Quaternary geology of the upper Cook Inlet area and Matanuska River Valley, in Péwé, T.L., ed., Guidebook for Field Conference F, VIIth Congress, International Association for Quaternary Research: Central and south-central Alaska: Fairbanks, Alaska Division of Geological and Geophysical Surveys, reprinted 1977, p. 114-141.
- Kromer, B., Rhein, M., Bruns, M., Schoch-Fischer, H., Munnich, K.O., Stuiver, M., and Becker, B., 1986, Radiocarbon calibration data for the 6th and 8th millennia BC: Radiocarbon, v. 28, p. 954-960.
- Linick, T.W., Suess, H.E., and Becker, B., 1985, La Jolla measurements of radiocarbon in South German Oak tree-ring chronologies: Radiocarbon, v. 27, p. 20-32.
- Linick, T.W., Long, A., Damon, P.E., and Ferguson, C.W., 1986, High-precision radiocarbon dating of Bristlecone Pine from 6554 to 5350 BC: Radiocarbon, v. 28, p. 943-953.
- McCann, W.R., Nishenko, S.P., Sykes, L.R., and Krause, J., 1979, Seismic gaps and plate tectonics: Seismic potential for major plate boundaries: Pure and Applied Geophysics, v. 117, p. 1082-1147.
- Ovenshine, A.T., Lawson, D.E., and Bartsch-Winkler, S.R., 1976, The Placer River Silt--An intertidal deposit caused by the 1964 Alaska earthquake: Journal of Research of the U.S. Geological Survey, v. 4, no. 2, p. 151-162.
- Pearson, G.W., and Stuiver, M., 1986, High-precision calibration of the radiocarbon time scale, 500-2500 BC: Radiocarbon, v. 28, p. 839-862.
- Pearson, G.W., Pilcher, J.R., Baille, M.G., Corbett, D.M., and Qua, F., 1986, High-precision ^{14}C measurement of Irish Oaks to show the natural ^{14}C variations from AD 1840-5210 BC: Radiocarbon, v. 28, p. 911-934.
- Plafker, George, 1969, Tectonics of the March 27, 1964, Alaska earthquake: U.S. Geological Survey Professional Paper 543-I, 74 p., scales 1:2,000,000 and 1:500,000, 2 sheets.
- _____ 1990, Regional vertical tectonic displacement of shorelines in south-central Alaska during and between great earthquakes (abs.), in Environmental disasters: The nature of recovery, Symposium, March 21, 1990, Corvallis, Oregon, Proceedings: Northwest Scientific Association.
- Plafker, George, and Rubin, Meyer, 1967, Vertical tectonic displacements in south-central Alaska during and prior to the great 1964 earthquake: Journal of Geosciences, Osaka City University, v. 10, p. 53-66.
- _____ 1978, Uplift history and earthquake recurrence as deduced from marine terraces on Middleton Island, Alaska, in Proceedings of Conference VI, Methodology for Identifying Seismic Gaps and Soon-to-break Gaps, National Earthquake Hazards Reduction Program, 25-27 May, 1978: U.S. Geological Survey Open File Report 78-943, p. 687-721.

- Reger, R.D., and Updike, R.G., 1983, Upper Cook Inlet region and the Matanuska Valley, *in* Péwé, T.L., and Reger, R.D., eds., Guidebook to permafrost and Quaternary geology along the Richardson and Glenn Highways between Fairbanks and Anchorage, Alaska: Alaska Division of Geological and Geophysical Surveys Guidebook 1, p. 185-263, scale 1:250,000, 1 sheet.
- Stuiver, M., Kromer, B., Becker, B., and Ferguson, C.W., 1986, Radiocarbon age calibration back to 13,300 years BP and the ^{14}C age matching of the German Oak and U S Bristlecone Pine chronologies: *Radiocarbon*, v. 28, p. 969-979.
- Stuiver, M., and Pearson, G.W., 1986, High-precision calibration of radiocarbon time scale, AD 1950-500 BC: *Radiocarbon*, v. 28, p. 805-838.
- Stuiver, M., and Reimer, P.J., 1986, A computer program for radiocarbon age calibration: *Radiocarbon*, v. 28, p. 1022-1030.
- U.S. Department of Commerce, 1988, Tide tables 1989, west coast of North and South America: Rockville, Maryland, National Ocean Survey, 222 p.
- Ward, P.L., and Page, R.A., 1989, The Loma Prieta earthquake of October 17, 1989: *Earthquakes & Volcanoes*, v. 21, no. 6, p. 215-246.
- West, D.O., and McCrumb, D.R., 1988, Coastal uplift in Oregon and Washington and the nature of Cascadia subduction-zone tectonics: *Geology*, v. 16, no. 2, p. 169-172.

APPENDIX A

BOREHOLE LOGS

All values in inches.

TA1 - Girdwood

9/19/88

Attempted		Recovered		Total	Total	Sample Number	Notes
Top	Bottom	Top	Bottom	Att.	Rec.		
0	42	0	19	42	19	TA1-1	Sample compressed in fill
42	72	42	51	30	9	TA1-2	Head clogged with gravel
72	102	72	86	30	14	TA1-3	Fibric peat at base (sedge, sphagnum, occ. wood)
		86	88		2	TA1-3C	Catcher - all organics
102	132	102	122	30	20	TA1-4	Organics at top
		125	132		7	TA1-4B	Recovered with next sample; disturbed
132	162	132	162	30	30	TA1-5	Organics(?) at 157 in.
162	222	162	187	60	25	TA1-6A	
		187	217		30	TA1-6B	Twig(?) at 195
222	252	222	242	30	20	TA1-7	Sand flowed in, drilled out to retry
252	282	252	282	30	30	TA1-8	Mostly peat
282	342	282	309	60	27	TA1-9A	Organics 282-294, 307-309
		309	337		28	TA1-9B	Organics 309-315
		337	340		3	TA1-9C	Catcher sample
342	372	342	372	30	30	TA1-10	
372	409	372	381	37	9	TA1-11A	Possibly disturbed - part of interval lost on TA1-10 attempt
		381	409		28	TA1-11B	Rest of lost interval from TA1-10
409	462	409	419	53	10	TA1-12A	Recovered from TA1-11 attempt
		419	449		30	TA1-12B	Dense silty fine sand
		449	452		3	TA1-12C	Catcher sample
462	522	462	490	60	28	TA1-13A	No organics visible
		490	520		30	TA1-13B	No organics visible
		520	522		2	TA1-13C	Catcher - dense silty fine sand
522	582	522	541	60	19	TA1-14A	Disturbed - flowed when laid horiz. for disassembly; no organics
		541	571		30	TA1-14B	No organics visible
		571	573		2	TA1-14C	Catcher sample
582	642	582	609	60	27	TA1-15A	Pebbles at top, Macoma shells at bottom; vy fine sand
		609	639		30	TA1-15B	Macoma shells at top
		639	642		3	TA1-15C	Catcher sample; dense vy fine sand
642	672	642	669	30	27	TA1-16	Very hard; filled 30 in. liner - discard top 3 in. - slough
		669	672		3	TA1-16C	Catcher sample
672	702	672	698	30	26	TA1-17	Wet, flowing.
		698	701		3	TA1-17C	Catcher
702	762	702	732	60	30	TA1-18A	Wet, flowing.
		732	762		30	TA1-18B	No organics; gravel at bottom (till?); very hard drilling last 2 ft.
				-----	-----		
				762	664		87.1% recovered

BOREHOLE LOG
TAB - Portage

9/19/88

Attempted		Recovered		Total Att.	Total Rec.	Sample Number	Notes
Top	Bottom	Top	Bottom				
0	42	0	27	42	27	TAB-1	Surface organics (1 in.), rest clean med-coarse sand
		27	29		2	TAB-1C	Catcher sample
42	102	42	47	60	5	TAB-2A	Bagged disturbed upper part: fine sand w/some silt, organic smears
		47	77		30	TAB-2B	
		77	80		3	TAB-2C	Catcher sample
102	132	102	131	30	29	TAB-3	Clean gray medium sand
		131	132		1	TAB-3C	Catcher sample
132	162	132	159	30	27	TAB-4	Peat layers at 140-142 and 150-159
		159	162		3	TAB-4C	Catcher sample - sedge peat
162	222	162	183	60	21	TAB-5A	Silty fine sand to silty clay
		183	213		30	TAB-5B	Silty clay & clayey silt w/scattered organics
		213	218		5	TAB-5C	Catcher sample
222	252	222	248	30	26	TAB-6	Sedge peat, bottom 8 in.
		248	252		4	TAB-6C	Catcher sample - sedge peat
252	282	252	279	30	27	TAB-7	Sedge peat top 2 in., organic silt.
		279	282		3	TAB-7C	Catcher sample - Very fine sand w/some silt
282	312	282	309	30	27	TAB-8	Sedge peat 297-301
		309	312		3	TAB-8C	Catcher, very fine sand w/some silt
312	342	312	339	30	27	TAB-9	Vy fine sand w/some silt and scattered organics
		339	342		3	TAB-9C	Catcher sample
342	402	342	372	60	30	TAB-10A	Very fine sand w/some silt; sphagnum peat 368-372
		372	402		30	TAB-10B	Peat 372-374; rest silty vy fine sand (w/some clay to 382 in.)
402	462	402	432	60	30	TAB-11A	Very fine sand w/some silt
		432	462		30	TAB-11B	Vy fine sand, clayey silt, organic silt, thin peat 455-456
462	522	462	492	60	30	TAB-12A	Sandy silt, clayey silt, organic silt 477-478
		492	522		30	TAB-12B	Clayey silt, silty fine sand
522	582	522	552	60	30	TAB-13A	Clayey silt, silty clay, silty fine sand
		552	582		30	TAB-13B	Clayey silt & silty clay; organic silt 555-556
582	642	582	612	60	30	TAB-14A	Clayey silt, silty fine sand, silty clay; trace organics at 594, 605-609
		612	642		30	TAB-14B	Clayey silt to silty clay
				-----	-----		
				642	603		93.9% recovered

BOREHOLE LOG

2/22/89

KA1 - Palmer Hayflats, 3 mi west of Glenn Highway and 1/4 mi north of Matanuska River.

All values in inches.

Attempted		Recovered		Total Att.	Total Rec.	Sample Number	Notes
Top	Bottom	Top	Bottom				
0	18	8	12	18	4	KA1-1	0-8" Ice. Bagged; frozen peat and silt.
18	48	18	32	30	14	KA1-2	Gray organic silt 18-24"; brown peat 24-32".
		32	36		4	KA1-2C	Catcher sample; basal peat with small amount of underlying gray silt.
48	78	48	72	30	24	KA1-3	Brownish-gray organic-rich clayey silt 48-55"; gray clayey silt with organic-rich zones 55-72".
		72	75		3	KA1-3C	Catcher sample; gray clayey silt with some organics.
78	108	84	102	30	18	KA1-4	Gray silt to clayey silt; no organics, wet.
		102	105		3	KA1-4C	Catcher; gray clayey silt.
108	138	105	135	30	30	KA1-5	Gray clayey silt with some organics(?) near top.
		135	136		1	KA1-5C	Catcher; gray silt.
138	168	138	166	30	28	KA1-6	Gray clayey silt 138-144"; Brownish gray fine sandy(?) silt with organics 144-146"; gray to dark gray organic silt 146-147"; gray clayey silt 147-166".
		166	167		1	KA1-6C	Catcher; gray clayey silt.
168	198	168	194	30	26	KA1-7	Gray silt.
		194	196		2	KA1-7C	Catcher; gray silt.
198	228	198	225	30	27	KA1-8	Gray silt.
		225	228		3	KA1-8C	Catcher; gray silt with some organics.
228	258	228	258	30	30	KA1-9	Massive gray silt.
						KA1-9C	Catcher (extra); gray silt.
258	288	258	288	30	30	KA1-10	Massive gray silt; some organic smears near bottom.
		288	292		4	KA1-10C	Catcher; gray silt.
288	348	292	317	60	25	KA1-11	Massive gray silt with minor organics in top 7'.
		317	347		30	KA1-12	Massive gray silt with minor organic smears.
		347	349		2	KA1-12C	Catcher; gray silt.
348	408	349	368	60	19	KA1-13	Massive gray silt with some black organic smears.
		368	398		30	KA1-14	Massive gray silt with abundant black organic smears.
		398	401		3	KA1-14C	Catcher; gray silt.
							Ended drilling at 34 ft; no more peat layers. Moved drill to 4 ft south of KA1 to re-sample 3-4 ft zone:
24	54	25	53	30	28	KA1B-1	Grayish brown silt-rich peat 25-30"; brownish gray silt with some organics 30-32"; grayish brown peat with some silt 32-36; gray silt with some black organic smears 36-53".
		53	55		2	KA1B-1C	Catcher; gray silt with minor organics.
				----- 438	----- 391		89.3% recovered

BOREHOLE LOG
KA2 - Palmer Hay Flats

2/15/89

All values in inches.

Attempted		Recovered		Total	Total	Sample Number	Notes
Top	Bottom	Top	Bottom	Att.	Rec.		
						--	-12 to 0' ice.
0	18	0	12	18	12	KA2-1	Bagged - frozen ice-rich peat.
18	36	18	34	18	16	KA2-2	Peat 18-21'; firm gray silt 21-34'
36	66	36	66	30	30	KA2-3	Massive gray silt to fine sand.
66	96	66	90	30	24	KA2-4	Massive fine sand; upper 6" probably disturbed - liquefies easily.
		90	93		3	KA2-4C	Catcher sample; gray fine sand with abundant mica, little or no silt; fluvial or tidal-channel bedload.
96	126	96	116	30	20	KA2-5	Massive gray fine sand; firm, liquefiable.
		118	119		3	KA2-5C	Catcher; firm gray fine sand.
126	156	126	147	30	21	KA2-6	Firm gray silty fine sand and silt with some rust coloring; no organics.
		147	149		2	KA2-6C	Catcher; gray silt.
156	216	156	180	60	24	KA2-7	Massive gray silt; organic streaks 174-177'.
		180	210		30	KA2-8	Massive gray silt; organic streaks 202-205'.
		210	212		2	KA2-8C	Catcher; gray silt.
216	246	216	246	30	30	KA2-9	Gray silt; firm, massive, no organics.
246	276	246	276	30	30	KA2-10	Firm, massive gray silt; black organic(?) smears 269-274'.
276	306	276	303	30	27	KA2-11	Very firm, massive gray silt; some black organic smears.
		303	306		3	KA2-11C	Catcher; very firm, massive gray silt.
306	336	306	331	30	25	KA2-12	Firm, massive gray silt; very hard drilling. Some black organic smears.
		331	334		3	KA2-12C	Catcher; same as above.
336	366	336	352	30	16	KA2-13	Very firm, massive gray silt with minor organic smears.
		352	355		3	KA2-13C	Catcher; same as above.
366	396	366	378	30	12	KA2-14	Very firm, massive gray silt.
		378	381		3	KA2-14C	Catcher; same as above.

Ended drilling at 33 ft. Silt too hard; over-torquing drill. Cannot drill further without damaging auger or sampler.

396 339

85.6% recovered

BOREHOLE LOG
KA3 - Palmer Hay Flats

2/15-2/16/89

Site about 2.5 miles WSW of BP well site.

Surface flat with tussocks; vegetation mostly grasses and dwarf birch, max. 2 ft high.

All values in inches.

Attempted		Recovered		Total	Total	Sample	Notes
Top	Bottom	Top	Bottom	Att.	Rec.		
						--	Augered through surface and frozen soil to 20" below ice surface (8" below top of soil. -12 to 0" Surface ice.
0	12	4	8	12	4	KA3-1	Bagged - frozen basal surface peat over gray silt.
12	36	12	15	24	3	KA3-2	Bagged - frozen gray silt, plugged sampler.
		15	26		11	KA3-3	Recovered lost portion; gray sandy silt, no organics.
36	66	33	63	30	30	KA3-4	Massive gray fine sandy silt; no organics.
		63	66		3	KA3-4C	Catcher sample - same as above.
66	96	66	96	30	30	KA3-5	Massive gray fine sandy silt; wet at top, moist at bottom.
96	156	96	110	60	14	KA3-6	Same.
		110	140		30	KA3-7	Same.
		140	142		2	KA3-7C	Catcher; same.
156	186	156	182	30	26	KA3-8	Same.
		182	185		3	KA3-8C	Same; one small wood fragment.
186	216	186	214	30	28	KA3-9	Gray silty fine sand, possibly fluvial.
		214	216		2	KA3-9C	Gray silty fine sand; scattered small wood fragments.
216	246	216	241	30	25	KA3-10	Saturated gray fine sand, mostly liquefied (section disturbed).
246	276	241	268	30	27	KA3-11	Same.
		268	271		3	KA3-11C	Gray fine sand.
				-----	-----		Stopped drilling at 23 ft; severe flow-in. Not able to sample without wet sand flowing in. Drilling ended at 2:00 pm, 2/16/89.
				276	241		87.4% recovered

BOREHOLE LOG
KA4 - Goose Bay

1/19 - 1/20/89

All values in inches.

Attempted		Recovered		Total	Total	Sample	Notes
Top	Bottom	Top	Bottom	Att.	Rec.		
						--	-12 to 0' Ice.
0	36	0	13	36	13	KA4-1	Frozen silt with peat at base.
		13	18		5	KA4-1C	Catcher sample; brown fibrous peat (1964 soil?) No recovery 18-36"; Fibrous peat with large sticks. Sampler probably plugged with frozen sediment and pushed peat aside. Sticks may be roots of 1964 vegetation.
36	66	36	48	30	12	KA4-2	Discard top 2' of 14" core (fill-in), 36-40" brown peat, 40-48" silt.
		48	51		3	KA4-2C	Catcher; silt.
66	96	66	93	30	27	KA4-3	Gray silt, slightly varved, faint rhythmic bedding.
		93	96		3	KA4-3C	Catcher; gray silt with trace organics.
96	156	96	106	60	9	KA4-4	Silt, no organics.
		105	135		30	KA4-5	Massive silt, no organics.
		135	138		3	KA4-5C	Catcher; silt.
156	186	156	179	30	23	KA4-6	Silt, no organics.
		179	182		3	KA4-6C	Catcher; silt.
186	216	186	211	30	25	KA4-7	Silt; trace organics at 187". Dark gray 203-211".
		211	214		3	KA4-7C	Catcher; silt.
216	246	216	242	30	26	KA4-8	Silt; trace organics(?) at top.
		242	245		3	KA4-8C	Catcher; silt.
246	276	246	270	30	24	KA4-9	Firm silt; faint bedding. No organics.
		270	273		3	KA4-9C	Catcher; firm silt.
276	306	276	301	30	25	KA4-10	Very firm silt; no organics.
		301	304		3	KA4-10C	Catcher; very firm silt.
306	336	306	324	30	18	KA4-11	Very firm, dry silt; split liner from being packed so tight. No organics.
		324	328		2	KA4-11C	Catcher; very firm, dry silt, no organics.

Ended drilling at 326" (27'2") because no organics were encountered below upper 18 inches. Prepared to move to alternate site to north up bay.

336 263

78.3% recovered

BOREHOLE LOG 2/13/89
 KA4B - Goose Bay (~ 1/4 mi NE of KA4)

Attempted		Recovered		Total	Total	Sample	Notes
Top	Bottom	Top	Bottom	Att.	Rec.		
						--	-12 to 0' ice.
0	12	0	10	12	10	KA4B-1	Frozen organic silt with roots.
		10	12		2	KA4B-1C	Catcher sample - organic silt.
12	36	12	17	24	5	KA4B-2C	Only 5' recovered & bagged. Organic silt, frozen. Sampler must be plugged with frozen soil and pushing soft soil aside.
36	66	36	40	30	4	KA4B-3C	4' recovered and bagged.
66	96	66	84	30	18	KA4B-4	Brown peat & twigs 66-70. Steel gray sticky silt 70-84.
		84	87		3	KA4B-4C	Catcher - gray sticky silt with organics.
96	126	96	126	30	30	KA4B-5	Gray silt with scattered organics. Concentrated organics (peat lenses) at 97-98' and 121-122'.
126	156	126	156	30	30	KA4B-6	Gray silt with scattered organics, possibly varved. Peat(?) lense at 127-128'.
156	186	156	180	30	4	KA4B-6C	Bagged sample. Gray sticky silt with minor organics. Lost some sample that fell out while pulling up sampler.
		180	172		12	KA4B-7	Recovered some of lost sample; possibly disturbed. Stiff gray silt.
186	216	186	205	30	19	KA4B-8	Gray silt.
		205	207		2	KA4B-8C	Catcher sample; same.
216	246	216	246	30	30	KA4B-9	Gray silt.
246	276	246	276	30	30	KA4B-10	Gray silt.
276	306	278	306	30	30	KA4B-11	Gray silt.

Stopped at 306' (25.5') because no more organics encountered and silt getting very firm and difficult to drill. See KA4C for re-sample of upper 10 ft.

 306 229 74.8% recovered

BOREHOLE LOG 2/13/89
 KA4C - Goose Bay (2' south of KA4B to re-sample upper 10 ft)

Attempted		Recovered		Total	Total	Sample	Notes
Top	Bottom	Top	Bottom	Att.	Rec.		
						--	-12 to 0' ice.
0	36			36	0	--	Recovered only dirty ice shavings (about 6") - no sample.
36	66	36	46	30	10	KA4C-1C	Bagged; all peat.
66	96	66	77	30	11	KA4C-2	Pushed in ahead of auger. Brown peat 66-70"; gray silt with scattered organics 70-77'.
96	126	96	126	30	30	KA4C-3	Upper 9' peat rich, may have fallen from upper zone (discard). 96-105 Questionable (peat). 105-107 Gray silt with organics. 107-110 Organic silt or peat. 110-121 Gray silt. 121-123 Organic silt. 123-126 Gray silt.

 126 51 40.6% recovered

BOREHOLE LOG
KA5 - Palmer Hay Flats

2/17/89

Site east of Well Site Road about 1/4 mi west of Weir Lake. Vegetation grasses, dwarf birch(?), willows, and scattered spruce. Ice & frozen vegetation to 2 ft; water and vegetation/roots to 7 ft from surface.

All values in inches.

Attempted Top	Attempted Bottom	Recovered Top	Recovered Bottom	Total Att.	Total Rec.	Sample Number	Notes
							-24 to 0' Surface ice and frozen organics above vegetation mat.
48	78	56	75	30	19	KA5-1	0 to 48' Not sampled; water and vegetation. 56 to 61' Basal brown peat. 61 to 65' Gray silt with minor organics 65 to 67' Organic silt. 67 to 71' Gray silt with some organics. 71 to 72' Organic silt. 72 to 75' Gray silt.
		75	78		3	KA5-1C	Catcher sample; gray silt with minor organics.
78	108	78	100	30	22	KA5-2	Gray silt with faint organic-rich zones.
		100	103		3	KA5-2C	Catcher; gray silt.
		103	108		5	KA5-3	Recovered lost zone; gray silt or silty fine sand.
108	138	108	138	30	30	KA5-4	Massive gray silt; no organics.
		138	141		3	KA5-4C	Catcher; gray silt.
138	198	141	159	60	18	KA5-5	Massive gray silt.
		159	189		30	KA5-6	Massive gray silt.
		189	192		3	KA5-6C	Catcher; same.
		192	198		6	KA5-7	Bagged; recovered from previous interval.
198	228	198	228	30	28	KA5-8	Massive gray silt; no organics.
		226	228		2	KA5-8C	Catcher; gray silt.
228	288	228	240	60	12	KA5-9	Massive gray silt.
		240	270		30	KA5-10	Massive gray silt.
		270	273		3	KA5-10C	Catcher; same.
288	318	288	311	30	23	KA5-11	Massive gray silt.
		311	314		3	KA5-11C	Catcher; same.
318	348	318	324	30	6	KA5-12	Firm gray silt; plugging sampler (sampled with catcher fingers; without fingers, samples slide out).
348	378	348	370	30	22	KA5-13	Massive gray clayey silt.
		370	373		3	KA5-13C	Catcher; same.
				-----	-----		
				330	274		83.0% recovered

BOREHOLE LOG

2/22-2/23/89

KA6 - Palmer Hay Flats about 200 yd west of pipeline and 1 mi north of Matanuska River.

All values in inches.

Attempted		Recovered		Total	Total	Sample	Notes
Top	Bottom	Top	Bottom	Att.	Rec.		
							0-6" ice. 6-26" Gray silt with organics; not recovered.
23	26	23	26	3	3	KA6-1	Bagged; base of frozen silt.
26	48	26	41	22	15	KA6-2	Dark brownish gray peat mixed with silt 26-29"; gray silt with organics 29-36; brown peat with silt lenses 36-41".
48	78	48	62	30	14	KA6-3	Gray silt 48-52"; grayish brown peat 52-55"; gray silt 55-62".
		62	85		3	KA6-3C	Catcher sample; gray silt.
78	108	78	96	30	18	KA6-4	Gray silt.
		96	98		2	KA6-4C	Catcher; gray silt.
108	138	108	119	30	11	KA6-5	Gray silt with black organic smears.
		119	131		12	KA6-6	Gray silt.
138	168	138	167	30	29	KA6-7	Massive gray silt.
						KA6-8	Extra sample; may be flow-in(?) or from higher level; massive gray silt.
168	198	168	198	30	30	KA6-9	Massive gray clayey silt.
						KA6-10	Extra sample, possibly flow-in; gray silt with organics.
198	228	198	228	30	30	KA6-11	Gray silt.
228	258	228	253	30	25	KA6-12	Massive gray silt.
258	288	258	278	30	20	KA6-13	Massive gray clayey silt with some faint organic smears.
		278	281		3	KA6-13C	Catcher; gray clayey silt.
288	318	288	308	30	20	KA6-14	Gray clayey silt with dark gray organic(?) rich layers and smears near bottom.
		308	311		3	KA6-14C	Catcher; gray clayey silt.
318	348	318	345	30	27	KA6-15	Firm gray to olive gray clayey silt with numerous dark gray organic smears and layers, most concentrated in lower 2/3.
		345	348		3	KA6-15C	Catcher; firm gray clayey silt with 1/2" dark gray organic-rich layer.
348	378	348	378	30	30	KA6-16	Gray to olive gray clayey silt with dark gray organic smears and faint layering.
378	408	378	406	30	28	KA6-17	Gray clayey silt with faint dark gray organic smears near top.
		406	408		2	KA6-17C	Catcher; gray clayey silt.
408	438	408	438	30	30	KA6-18	Firm gray clayey silt with minor dark gray organic smears.
438	468	438	468	30	30	KA6-19	Gray clayey silt, not as firm as above.
468	528	468	496	60	28	KA6-20	Massive gray silt.
		496	526		30	KA6-21	Gray silt to fine-medium sand, very firm; split sample tube due to pressure (taped with electrical tape).
		526	528		2	KA6-21C	Firm gray silty fine sand.
							11 ft of flow-in occurred at this depth; not possible to drill any more without flow-in; stopped drilling at 44 ft.
				505	448		88.7% recovered

APPENDIX B

SAMPLE LOG

Descriptions of core samples upon opening in the laboratory.

BOREHOLE TA1, GIRDWOOD - 60°56'39"N, 149°10'35"W; surface elevation 20.7 ft MSL.

<u>SAMPLE</u>	<u>DEPTH (ft)</u>	<u>DESCRIPTION</u>
TA1-1	0-1.55	Silty, gravelly sand, very poorly sorted, nonstratified (partially disturbed during splitting because of large pebbles). Brownish gray to grayish brown. Fill.
	1.55-3.5	Not recovered.
TA1-2	3.5-3.75	Brownish gray, poorly sorted silty, gravelly sand.
	3.75-3.95	Dark brown to brownish gray silty, gravelly sand with abundant organics, including disseminated charcoal, rootlets, and one wood fragment.
	3.95-4.2	Gray to brownish gray, poorly sorted silty, sandy gravel; pebbles to 1 in. Entire sample (TA1-2) probably fill.
	4.2-6.0	Not recovered.
TA1-3	6.0-6.25	Gray, poorly sorted, silty, gravelly sand.
	6.25-6.35	Gray silty fine sand with some organics (sedge blades).
	6.35-6.5	Brown sedge peat. <u>Spl TA1-3-6.4</u> at top of peat layer, 6.35-6.45 (C14).
	6.5-7.2	Dark brown woody peat with abundant roots and twigs. Upper contact sharp along dark band.
TA1-3C	7.2-7.4	Same as 6.5-7.2 (catcher sample; bagged).
TA1B-1		(attempted resample of missed interval 7.4-8.5 ft depth, 5 ft SE of TA1)
	6.0-6.75	Gray very poorly sorted silt, sand, and gravel. Large rounded pebble at top, 0.2 in. diameter. Pebbles very angular to subrounded, mostly angular. Sharp lower contact. Possibly fill.
	6.75-7.4	Brown to dark brown fibrous peat with abundant roots and twigs. Lighter colored grassy layer in top 0.1 in. (sedge peat).
TA1B-1C	7.4-7.6	Peat as above (catcher sample; bagged).
	7.6-8.5	Not recovered.
TA1-4A	8.5-8.85	Dark brown woody peat; abundant twigs and roots. <u>Spl TA1-4A-8.8</u> basal peat, 8.75-8.85 (C14).
	8.85-10.1	Interlayered gray to brownish-gray silty fine sand and sandy silt. Finely laminated between 9.2 and bottom. Laminations are horizontal and less than 1/8 in. thick. A brownish rind exists about 1/4 in. thick along sample margins, possibly from iron-rich water penetrating along edges during sampling. Laminations are visible in rind, so rind is not due to disturbance during sampling.
	10.1-10.5	Not recovered.
TA1-4B	10.5-11.0	Massive gray silt; one 1/2 in. diameter twig, probably transported.
TA1-5	11.0-11.4	Brownish-gray silt and fine sand; disturbed, probably by sampling.
	11.4-11.8	Gray silt and fine sand with contorted, thin, indistinct laminae.

	11.8-12.75	Gray to brownish gray silt, possibly clay-rich, with scattered twigs and sedge blades. Faint, thin horizontal laminae 1-4 mm thick. Twig, 1/2 in. wide, at 12.1; probably transported. Sharp lower contact.
	12.75-13.1	Dark brown to light brown sedge peat. Gradational lower contact. <u>Spl TA1-5-12.8</u> at top of peat layer, 12.75-12.85 (C14).
	13.1-13.5	Gray and stained (mottled) light brown clay and silt with scattered organics. No visible bedding.
TA1-6A	13.5-13.9	Gray to brownish gray silt/fine sand with some iron staining along edges. Trace organics.
	13.9-15.4	Gray to brownish gray silt/fine sand with distinct laminations 1-4 mm, generally horizontal and undisturbed. Scattered roots and twigs. Flat piece of wood 1 in. x 1 in. x 1/4 in. at 15.2, partially charcoal; probably transported.
TA1-6B	15.4-15.8	Disturbed zone (from sampling) of gray silt/fine sand with some iron staining around scattered organics (roots or sedge blades).
	15.8-17.1	Laminated to thin-bedded gray silt and fine sand with scattered sedge blades or roots. Iron stained around organics.
	17.1-17.45	Laminated gray silt and clayey silt. Trace organics. <u>Spl TA1-6B-17.1</u> at 17.0-17.2 (GS).
	17.45-17.75	Dark brownish gray pebbly medium to coarse sand; trace silt. No visible bedding. Pebbles subangular to well rounded.
TA1-6C	17.75-17.9	Pebbly coarse sand. Angular to subrounded pebbles to 1/2 in. diameter (catcher sample; bagged).
	17.9-18.5	Not recovered.
TA1-7	18.5-18.8	Gray medium to coarse clean sand, coarsening downward; silt lense at 18.7. A few twigs or roots at top surface; 1/4-in. twig at 18.65, probably transported.
	18.8-19.3	Gray pebbly coarse sand with angular to rounded pebbles to 3/4 in. Negligible silt, no visible bedding. <u>Spl TA1-7-19.0</u> at 18.9-19.2 (GS).
	19.3-19.6	Gray medium to coarse sand with some small pebbles and negligible silt, coarsening downward.
	19.6-19.65	Uniform gray clayey silt.
	19.65-20.2	Dark brown woody or sedge peat. Twigs up to 1/4 in. diameter. <u>Spl TA1-7-19.7</u> at 19.65-19.75 (C14).
TA1-7C	20.2-20.4	Peat as above (catcher sample; bagged).
	20.4-21.0	Not recovered.
TA1-8	21.0-21.1	Gray silt, sand, pebbles, and organics. Could be slough from drill stem.
	21.1-22.15	Dark brown layered woody or sedge peat with a few thin lenses of gray silt. <u>Spl TA1-8-21.2</u> of surface peat, 21.1-21.25 (C14).
	22.15-22.5	Gray clayey silt with some twigs and roots.
	22.5-23.5	Dark brown woody peat with minor thin silt layers near top. Lighter brown in upper 3 in. <u>Spl TA1-8-22.6</u> of surface peat, 22.5-22.7 (C14).
TA1-9A	23.5-23.6	Mixed silt, sand, and organics; probably disturbed during sampling.
	23.6-24.3	Layered dark brown sedge peat and gray silt. Possibly gradational base of lower peat in TA1-8.

	24.3-25.4	Massive gray clayey silt mottled with dark gray organic smears, but organics are negligible. <u>Spl TA1-9A-25.3</u> at 25.2-25.4 (GS).
	25.4-25.6	Dark brown sedge and woody peat. <u>Spl TA1-9A-25.5</u> at 25.4-25.6 (C14).
TA1-9B	25.6-25.9	Dark brown woody peat grading downward to brownish gray clayey silt, faintly layered (base of peat unit at bottom of TA1-9A).
	25.9-27.85	Brownish gray to gray clayey silt with scattered organics (roots and sedge blades). Very faint thin layering. Lower part mottled with dark gray organic smears. <u>Spl TA1-9B-26.3</u> at 26.2-26.4 (GS).
TA1-9C	27.85-28.1	Gray clayey silt (catcher sample; bagged).
	28.1-28.5	Not recovered.
TA1-10	28.5-30.95	Brownish gray to gray clayey silt. Very faint thin layering. Iron staining in lower portion concentrated around minor organics. Some dark gray mottling, probably organic smears.
TA1-11A	31.0-31.65	Gray clayey silt with dark gray organic smears. Trace sedge blades and minor iron staining along edges.
TA1-11B	31.65-33.8	Gray to brownish gray clayey silt with trace organics, minor iron staining, and dark gray organic smears. Grades to sandier near bottom; minor small pebbles in lower 3 in. <u>Spl TA1-11B-33.0</u> at 32.9-33.1 (GS).
TA1-12A	33.8-34.55	Brownish gray to gray silt and fine sand; no visible bedding, possibly disturbed.
TA1-12B	34.55-37.0	Brownish gray fine sandy silt and clayey silt with pebbly zones. Mottled with dark silty zones. Interlayered dark gray silt and brownish gray silty fine sand 36.3-36.8, layers 1/2 to 1 in. thick. All but 36.3-36.8 may be disturbed. No organics. Single angular pebble at 34.8. Pebbly (angular to rounded) and sandy zone 35.4-35.7. Two-inch rounded pebble at 36.8. <u>Spl TA1-12B-36.5</u> at 36.4-36.6; layered sandy silt and clayey silt (GS).
TA1-12C	37.0-37.25	Coarse silt or silty fine sand (catcher sample; bagged).
	37.25-38.5	Not recovered.
TA1-13A	38.5-40.75	Brownish gray to gray silt and fine sand with minor scattered pebbles. Most pebbles subrounded to well rounded; max. diameter about 1 in. Large (2-1/2 in.) very angular pebble at 40.5. No visible bedding. Much of this sample may be disturbed, probably during transport.
TA1-13B	40.75-41.5	Disturbed layer of brownish gray silty fine sand.
	41.5-43.25	Interlayered light brown silty fine sand and gray fine sandy silt. Some thin bedding and lamination. A few pebbles, including one subrounded pebble 1 in. diameter at 42.6. No organics. <u>Spl TA1-13B-41.8</u> at 41.7-41.9, brown sandy silt (GS).
TA1-13C	43.25-43.4	Dense silty fine sand (catcher sample; bagged).
	43.4-43.5	Not recovered.
TA1-14A	43.5-44.75	Dark gray uniform silty fine sand. No visible bedding. Trace pebbles (1/2-in. angular pebble at 44.6). No organics. Could be disturbed (according to drilling log).
TA1-14B	44.75-45.3	Gray to brownish gray silty fine sand. Slight hint of thin bedding. One rounded pebble 1-3/4 in. long at 44.8.
	45.3-45.4	Gray clayey silt, uniform.

	45.4-46.7	Grayish brown to gray silt with distinct horizontal layering in lower 0.3 ft. Two pebbles about 2 in. long at 45.9 and 46.2. Slight grading to medium sand at base. <u>Spl TA1-14B-45.6</u> at 45.5-45.7 (GS).
	46.7-47.2	Gray to brownish gray silty fine sand. No visible bedding. Dark gray zone at 46.9-47.1 may be organic-rich (disseminated).
TA1-14C	47.2-47.4	Silty fine sand or coarse silt (catcher sample; bagged).
	47.4-48.5	Not recovered.
TA1-15A	48.5-50.4	Brownish gray to gray uniform silty fine sand. Brownish (iron stained) 48.5-48.8. Minor pebbles (rounded, up to 1/2 in.) at 48.6 and 50.0. Trace shell fragments (48.65, 49.9).
TA1-15B	50.4-52.75	Gray to brownish gray uniform silty fine sand. Trace shell fragments (e.g. 50.7). Several angular to rounded pebbles to 1/2 in. at 52.6.
TA1-15C	52.75-53.0	Dense, very fine sand (catcher sample; bagged).
	53.0-53.5	Not recovered.
TA1-16	53.5-53.9	Slough from drill stem.
	53.9-55.9	Brownish gray to gray uniform silty fine sand or sandy silt. Very faint layering. One-inch angular pebble at 54.0 (probably ice-rafted). Dark gray layers at 55.2-55.4, organic-rich(?). <u>Spl TA1-17-55.3</u> at 55.2-55.4, thin laminated charcoal in silty fine sand (C14).
TA1-16C	55.9-56.0	Silty fine sand or sandy silt (catcher sample; bagged).
TA1-17	56.0-58.0	Brownish gray to gray uniform fine sandy silt. No visible bedding and no organics. Trace scattered small pebbles. Brownish color apparently from iron staining. <u>Spl TA1-17-57.0</u> at 56.9-57.1 (GS).
TA1-17C	58.0-58.25	Silty fine sand or sandy silt (catcher sample; bagged).
	58.25-58.5	Not recovered.
TA1-18A	58.5-60.6	Brownish gray to gray uniform silty fine sand with trace scattered small pebbles (probably ice-rafted). No visible bedding or organics. Brownish color from iron staining. Rounded pebble 3/4-inch at 60.4.
TA1-18B	60.6-62.3	Brownish gray to gray uniform fine sandy silt with trace scattered small pebbles (one well-rounded pebble 1/2-in. dia.) at 61.95. No visible bedding. Trace organics (dark gray smear) at 61.5. Brownish color from iron staining. <u>Spl TA1-18B-61.0</u> at 60.9-61.1 (GS).
	62.3-62.4	Thin layer of pebbly coarse sand.
	62.4-62.7	Gray uniform silty fine sand. No visible bedding or organics.
	62.7-63.05	Gray pebbly, silty fine sand; 2-in. pebble at 63.0.
TA1-18C	63.05-63.3	Gravel (catcher sample; bagged).

Maximum depth sampled 63.3 ft.

BOREHOLE TA8, PORTAGE- 60°48'54"N, 148°57'57"W; surface elevation 23.5 ft MSL.

<u>SAMPLE</u>	<u>DEPTH (ft)</u>	<u>DESCRIPTION</u>
TA8-1	0-0.3	Gray fine sand mixed with surface organics (leaves, twigs, roots).
	0.3-1.8	Gray to brownish gray, layered silty very fine sand to clean fine or medium sand. Four layers 0.2 to 0.6 ft thick, reverse graded (fine at bottom to medium at top). Scattered roots and twigs throughout. Mica rich 1.0-1.6.

	1.8-2.2	Gray uniform fine to medium sand, clean. No bedding; trace organics. Probably fluvial.
TA8-1C	2.2-2.4	Fine to medium sand with trace silt (catcher sample; bagged).
TA8B-1		(From adjacent borehole to resample missed interval 2.2-3.9; 6 ft NE of TA8)
	1.4-1.6	Disturbed. Gray fine to medium sand with some organics.
	1.6-3.1	Gray to brownish gray, layered silty very fine sand to clean fine or medium sand. Three layers to 0.6-in. thick, reverse graded as in TA8-1, 0.3-1.8. Trace organics.
TA8B-1C	3.1-3.2	Clean medium sand (catcher sample; bagged).
	3.2-3.5	Not recovered.
TA8-2A	3.5-3.9	Fine sand with some silt and organic smears (bagged).
TA8-2B	3.9-4.85	Gray, uniform, fine to medium sand with minor silt. No visible layering. Trace organics. <u>Spl TA8-2B-4.7</u> at 4.6-4.8 (GS).
	4.85-5.1	Interlayered brownish gray to gray silt, silty fine sand, and organics (probably sedge). <u>Spl TA8-2B-4.9</u> at 4.85-5.1 (C14); sedge blades, roots, and rhizomes(?), probably in place at time of deposition.
	5.1-6.3	Gray silt and silty fine sand with faint layering and scattered minor organics.
TA8-2C	6.3-6.55	Silt or silty fine sand (catcher sample; bagged).
TA8B-2		(From adjacent borehole to sample missed interval 6.3-8.4 in TA8; 6 ft NE of TA8)
	6.0-6.4	Probably disturbed, gray, uniform, clean fine sand; probably fluvial.
	6.4-6.9	Gray silt and silty fine sand with faint layering. Minor iron staining. Trace organics.
	6.9-7.2	Brownish gray to gray layered silt, silty fine sand, and organics (sedge). Iron stained in organic zones. Fluvial(?). <u>Spl TA8B-2-7.0</u> at 6.9-7.1 (C14).
	7.2-8.1	Brownish gray to gray silt, silty fine sand, and clean fine sand. Distinct layering (silt layer with sharp contacts at 7.9-8.0). Negligible organics.
TA8B-2C	8.1-8.3	Clean fine sand (catcher sample; bagged).
	8.3-8.5	Not recovered.
TA8-3	8.5-10.7	Gray silty fine sand and clean fine sand with some silt interlayers 10.0-10.7. Faint iron staining. No organics. <u>Spl TA8-3-9.5</u> at 9.4-9.6 (GS).
TA8-3C	10.7-11.0	Fine sand with trace silt (catcher sample; bagged).
TA8-4	11.0-11.85	Gray silt, silty fine sand, and clean fine sand in contorted zones; probably disturbed during sampling. Some evidence of lamination at 11.4. Negligible organics. Minor iron staining.
	11.85-12.6	Gray to brownish gray clayey silt and fine sandy silt. Disseminated organics 12.1-12.3. Scattered sedge blades. <u>Spl TA8-4-12.5</u> at 12.4-12.6 (GS).
	12.6-13.25	Brown to dark brown sedge and woody peat. Increasing silt toward base. Sharp upper contact. <u>Spl TA8-4-12.7</u> at 12.6-12.75 (C14).
	13.25-13.4	Brownish gray organic-rich silt.

TA8-4C	13.4-13.5	Sedge peat (catcher sample; bagged). <u>Spl TA8-4C-13.4</u> (C14).
TA8-5A	13.5-13.6	Gray fine sand; probably not in place (sloughed or fill-in). Lower contact distorted.
	13.6-15.2	Gray silt with abundant scattered organics (sedge blades, roots).
TA8-5B	15.2-17.5	Gray to brownish gray (minor) laminated silt and clayey silt. Horizontal laminations 1-5 mm. Trace scattered organics (sedge blades). <u>Spl TA8-5B-16.7</u> at 16.6-16.8 (GS).
TA8-5C	17.5-18.1	Fine sand with some silt and clay (catcher sample; bagged).
	18.1-18.5	Not recovered.
TA8-6	18.5-19.3	Disturbed gray clayey silt and silty fine sand. Contorted laminae. No organics.
	19.3-20.2	Gray to brownish gray (at base) laminated silt and clayey silt. Minor organics near base. Lower contact gradational. <u>Spl TA8-6-20.0</u> at 19.9-20.1 (GS).
	20.2-20.9	Woody and sedge peat, mixed with silt in upper 0.1 ft. <u>Spl TA8-6-20.3</u> at 20.2-20.35 (C14).
TA8-6C	20.9-21.0	Sedge peat (catcher sample; bagged).
TA8-7	21.0-21.4	Probably fill-in (not in place). Gray silt with chunks of peat.
	21.4-21.7	Dark brown sedge(?) peat grading downward to organic-rich silt. Probably the base of peat layer that starts at 20.2. Sampled to confirm. <u>Spl TA8-7-21.5</u> at 21.4-21.5 (C14).
	21.7-23.3	Gray clayey silt and silty fine sand with minor organics near top. Faint laminations near base.
TA8-7C	23.3-23.5	Very fine sand with some silt and trace clay (catcher sample; bagged).
TA8-8	23.5-24.75	Uniform gray silt or fine sandy silt with minor clay. Very faint laminations near base. Trace organics. <u>Spl TA8-8-24.6</u> at 24.5-24.7 (GS).
	24.75-25.0	Dark brown woody/sedge peat. Sharp upper contact, fairly sharp lower contact. <u>Spl TA8-8-24.8</u> at 24.75-24.85 (C14).
	25.0-25.75	Uniform gray silt or silty fine sand. Very faint thin laminations. No organics.
TA8-8C	25.75-26.0	Very fine sand with some silt (catcher sample; bagged).
TA8-9	26.0-28.3	Gray to brownish gray silt and silty fine sand with scattered minor organics (sedge). Disturbed with some contorted laminae in upper 1.0 ft. Top 0.3 ft may be fill-in (see log). Contorted layers of sedge blades or roots at 27.7. Distinct thin lamination at 27.2-27.5 and 27.8-28.3.
TA8-9C	28.3-28.5	Very fine sand with some silt (catcher sample; bagged).
TA8-10A	28.5-30.15	Uniform gray silt or silty fine sand with negligible organics. No visible bedding.
	30.15-30.25	Gray to brownish gray thin-bedded to laminated silt and clayey silt. Sharp lower contact with peat. <u>Spl TA8-10A-30.2</u> at 30.1-30.25 (GS).
	30.25-30.6	Dark brown sedge and/or sphagnum peat. Compact and horizontally layered. <u>Spl TA8-10A-30.3</u> at 30.15-30.4 (C14).
TA8-10B	30.6-30.8	Dark brown sedge and/or sphagnum peat. Base of peat layer at 30.25-30.6 in TA8-10A. Sharp base.

	30.8-33.1	Gray to brownish gray (minor) silt and silty fine sand. Organic-rich 30.8-31.1. Scattered organics decrease downward. Distinct fine laminations (0.5-2 mm) at 31.7-33.1. Minor iron staining along edges and near organics.
TA8-10C	33.1-33.2	Very fine sand (catcher sample; bagged).
	33.2-33.5	Not recovered.
TA8-11A	33.5-35.65	Gray clayey silt, nearly uniform. No visible bedding except fine laminations at 33.5-33.7 (top). Rest is massive or disturbed. Minor iron staining along margins. Negligible organics. Faint dark organic smears. <u>Spl TA8-11A-35.0</u> at 34.9-35.1 (GS).
TA8-11B	35.65-35.95	Gray to brownish gray silt. No bedding, possibly disturbed. Minor iron staining.
	35.95-36.2	Gray to brownish gray silt interlayered with organics. Concentrated organic layer 1/4 in. thick at 36.15. Possibly a buried peat layer. <u>Spl TA8-11B-36.2</u> at 36.15-36.25 (C14).
	36.2-38.1	Gray to brownish gray silt with interlayered thin organic-rich layers, decreasing downward. <u>Spl TA8-11B-36.8</u> at 36.8-36.9 (C14) for sedimentation rate.
	38.1-38.5	Not recovered.
TA8-12A	38.5-41.0	Gray silt. No visible bedding except faint laminations in lower 1.0 ft. trace organics. Trace iron staining.
TA8-12B	41.0-41.4	Gray laminated silt and silty fine sand. No visible organics.
	41.4-41.55	Gray to brownish gray silt with thin layers of organic material. Possibly a buried peat layer. <u>Spl TA8-12B-41.5</u> at 41.4-41.55 (C14).
	41.55-43.5	Gray to brownish gray silt with scattered organics and very faint laminations in lower 1.0 ft. No distinct upper contact. Distorted organic zone (sedge?) at 42.3.
TA8-13A	43.5-46.0	Gray clayey silt with trace organics and minor iron staining. Faint lamination at 44.8 to 46.0. <u>Spl TA8-13A-45.5</u> at 45.4 to 45.6 (GS).
TA8-13B	46.0-48.5	Gray silt and/or clayey silt with trace organics and minor iron staining. Faint laminations throughout, most visible at 47.0 where iron staining accentuates coarser laminae.
TA8-14A	48.5-50.95	Gray to brownish gray silt and clayey silt. No visible bedding. Minor iron staining (49.9-50.1). Mottled with dark gray organic smears throughout, especially visible at 50.1 to 50.95. Probably not a buried peat layer, but could be useful for determining sedimentation rate. <u>Spl TA8-14A-50.5</u> at 50.4-50.6 (GS). <u>Spl TA8-14A-50.8</u> at 50.6-50.9 (C14).
TA8-14B	50.95-53.4	Gray to brownish gray silt; laminated to thin bedded, except upper 0.5 ft probably disturbed. Light gray thin partings (volcanic ash?) 1 to 3 mm thick at 52.0 and 52.7-58.5.
TA8-14C	53.4-53.7	Silty clay with some fine sand (catcher sample; bagged).

Maximum depth sampled 53.7 ft.

BOREHOLE KA1, PALMER HAY FLATS - 61°29'40"N, 149°18'40"W; surface elevation 20.3 ft MSL.

<u>SAMPLE</u>	<u>DEPTH [ft]</u>	<u>DESCRIPTION</u>
	0-0.7	Not sampled; mostly ice.
KA1-1	0.7-1.0	Frozen peat and silt; bagged.

	1.0-1.5	Not recovered.
KA1-2	1.5-1.65	Wet gray silt mixed with grass, twigs, and roots, probably disturbed. Appears to be silt layer between surface organics and underlying peat.
	1.65-2.5	Grayish brown to dark brown peat (wood, sedge, sphagnum?). Larger pieces at top. Could have roots from surface. Some gray silt mixed in top 0.2 in. <u>Spl KA1-2-1.7</u> at 1.65-1.8 (C14).
KA1-2C	2.5-2.8	Basal peat with small amount of underlying gray silt (catcher sample; bagged).
KA1B-1		(From adjacent borehole to sample missed interval at 2.5-4.0 in KA1; 4 ft south of KA1)
	2.1-2.3	Wet gray silt mixed with sedge, twigs, and roots. Probably equivalent to 1.5-1.65 in KA1-2.
	2.3-2.8	Brown peat (wood, sedge, sphagnum?). Could be 1964 peat. Equivalent to peat at 1.65-2.5 in KA1-2. Sampled at base to compare with age of KA1-2-1.7 and avoid possible contamination by roots. <u>Spl KA1B-1-2.7</u> at 2.65-2.8 (C14).
	2.8-4.3	Wet gray clayey silt with faint thin lamination at top (2.8-3.1) and trace organics. <u>Spl KA1B-1-3.1</u> at 3.0-3.2 (GS).
KA1B-1C	4.3-4.5	Gray clayey silt with minor organics (catcher sample; bagged; from adjacent borehole 4 ft south of KA1).
KA1-3	4.0-5.8	Wet gray clayey silt with very faint layering and some organics. Also some dark gray organic smears. <u>Spl KA1-3-5.7</u> at 5.6-5.8 (GS).
	5.8-6.0	Dark brown sedge(?) peat grading downward to brownish gray organic-rich silt. Sharp upper contact. Probably buried peat layer. <u>Spl KA1-3-5.85</u> at 5.8-5.9 (C14).
KA1-3C	6.0-6.25	Gray clayey silt with some organics (catcher sample; bagged).
	6.25-7.0	Not recovered.
KA1-4	7.0-8.5	Gray silt with no visible bedding. Minor organics at 7.7-7.9 (sedge), otherwise rare. Dark gray organic smears throughout.
KA1-4C	8.5-8.75	Gray clayey silt.
KA1-5	8.75-11.3	Uniform gray clayey silt with no visible bedding and negligible organics. <u>Spl KA1-5-10.0</u> at 9.9-10.1 (GS).
KA1-5C	11.3-11.5	Gray clayey silt (catcher sample; bagged)
KA1-6	11.5-11.7	Gray clayey silt with no visible bedding. Trace organics. <u>Spl KA1-6-11.6</u> at 11.5-11.7 (GS).
	11.7-12.0	Dark brown peat (sedge?) grading down to grayish brown organic-rich silt. Upper contact slightly disturbed but sharp. Lower contact gradational. Probably a buried peat layer. <u>Spl KA1-6-11.75</u> at 11.7-11.8 (C14).
	12.0-13.8	Uniform wet gray silt with minor organics near top and trace at bottom. Faint laminations visible at 13.6-13.8.
KA1-6C	13.8-14.0	Gray clayey silt (catcher sample; bagged).
KA1-7	14.0-16.1	Wet clayey silt with faint layering (15.0-16.0) and no organics.
KA1-7C	16.1-16.3	Gray silt (catcher sample; bagged).

KA1-8	16.3-18.65	Uniform wet gray clayey silt. Very faint layering 18.0-18.4. Trace organics at 16.7. Several pockets of sedge blades at 18.1-18.5. Subrounded pebble 3/8 in. diameter at 18.35, probably ice rafted. <u>Spl KA1-8-17.9</u> at 17.8-18.0 (GS). <u>Spl KA1-8-18.2</u> at 18.1-18.3 (C14) for sedimentation rate (sedge blades probably transported).
KA1-8C	18.65-19.0	Gray silt with some organics (catcher sample; bagged).
KA1-9	19.0-21.5	Uniform wet gray clayey silt. Faint layering (20.9-21.1). Pockets of organics (sedge?) at 19.25-19.5.
KA1-10	21.5-23.95	Wet gray silt with faint layering (22.5-23.5) and minor scattered organics (e.g., 22.1, 23.2)
KA1-10C	23.95-24.3	Gray silt (catcher sample; bagged).
KA1-11	24.3-26.4	Gray clayey silt with some dark gray organic smears and moderate layering (see 25.1-25.6). Very minor organic pieces scattered throughout. <u>Spl KA1-11-25.0</u> at 24.9-25.1 (GS).
KA1-12	26.4-28.1	Uniform gray silt with minor dark gray organic smears. Small pocket of organics (sedge) at 27.1 Very faint layering.
	28.1-28.15	Layered gray silt and fine sand. Sand layer 3 mm thick at 28.15 above organic-rich layer. Too thin to sample for grain size.
	28.15-28.3	Dark gray layered organic-rich silt. Buried peat layer? <u>Spl KA1-12-28.2</u> at 28.15-28.3 (C14).
	28.3-28.9	Gray silt with some dark gray organic smears. Very faint layering.
KA1-12C	28.9-29.1	Gray silt (catcher sample; bagged).
KA1-13	29.1-30.7	Gray silt with very faint layering. Banded dark gray organic smears at 30.4-30.6. Doubtful buried peat layer, but would give sedimentation rate. <u>Spl KA1-13-30.5</u> at 30.4-30.6 (C14).
KA1-14	30.7-33.2	Gray silt with dark banded organic smears at 30.7-31.1, 31.3-31.6, and 32.4-32.5. Otherwise very faint layering. The three dark layers may be buried peats, but doubtful.
KA1-14C	33.2-33.5	Gray silt (catcher sample; bagged).

Maximum depth sampled 33.5 ft.

BOREHOLE KA2, PALMER HAY FLATS - 61°30'50"N, 149°26'15"W; surface elevation 19.3 ft MSL.

<u>SAMPLE</u>	<u>DEPTH (ft)</u>	<u>DESCRIPTION</u>
KA2-1	0-1.0	Frozen ice-rich peat (bagged).
	1.0-1.5	Not recovered.
KA2-2	1.5-1.75	Dark brown peat; base of modern surface organic layer.
	1.75-2.8	Uniform wet gray silt. No visible bedding. Trace organics (one small pocket of sedge blades at 2.65).
	2.8-3.0	Not recovered.
KA2-3	3.0-3.85	Gray silt. No visible bedding or organics. Minor iron staining. <u>Spl KA2-3-3.5</u> at 3.4-3.6 (GS).
	3.85-5.35	Gray clayey silt with faint laminations, grading downward to interlayered clayey silt and silty fine sand. Trace organics, minor iron staining. <u>Spl KA2-3-4.0</u> at 3.9-4.1 (GS).
	5.35-5.6	Not recovered.

KA2-4	5.5-7.3	Wet, gray silt, partially liquefied. Very faint bedding, trace organics. <u>Spl KA2-4-7.0</u> at 6.9-7.1 (GS).
KA2-4C	7.3-7.6	Gray fine sand with abundant mica and little or no silt (catcher sample; bagged). May be fluvial.
	7.6-8.0	Not recovered.
KA2-5	8.0-9.7	Gray clayey silt with some laminations of silty fine sand (9.0-9.7). Trace organics, minor iron staining.
KA2-5C	9.7-10.0	Firm gray fine sand (catcher sample; bagged).
	10.0-10.5	Not recovered.
KA2-6	10.5-12.2	Gray silt with iron staining along laminations. No visible layering. Possible volcanic-ash parting at 12.1 (very light gray silt-size material).
KA2-6C	12.2-12.4	Gray silt (catcher sample; bagged).
	12.4-13.0	Not recovered.
KA2-7	13.0-15.0	Gray clayey silt with no visible bedding. Minor iron staining. Trace scattered organics. Iron staining produces mottling, apparently associated with organics.
KA2-8	15.0-17.5	Gray clayey silt. Very faint bedding. Minor iron staining (mottled) at 15.0-16.4. Trace organics.
KA2-8C	17.5-17.7	Gray clayey silt (catcher sample; bagged).
	17.7-18.0	Not recovered.
KA2-9	18.0-20.5	Gray clayey silt. Distinct lamination and thin bedding at 19.5-20.3. No visible organics. Trace iron staining. <u>Spl KA2-9-20.0</u> at 19.9-20.1 (GS).
KA2-10	20.5-22.3	Gray clayey silt with faint layering. Desiccation crack longitudinally along entire sample. Several lateral cracks. No visible organics; no iron staining.
	22.3-23.0	Dark gray organic-rich clayey silt. Dark organic smears show crude bedding. Possible buried peat layer, or useful for age and sedimentation rate. <u>Spl KA2-10-22.4</u> at 22.3-22.45 (C14).
KA2-11	23.0-25.3	Gray clayey silt with no visible bedding. Mottled with small darker gray organic smears, but organics are negligible.
KA2-11C	25.3-25.5	Firm gray clayey silt (catcher sample; bagged).
KA2-12	25.5-27.5	Firm gray clayey silt with numerous dark gray organic smears in mottles and layers. Organic-rich zones (buried peat layers?) at 26.0-26.1, 26.7, and distinct layer at 27.05-27.15. Light brown fine-sand interbeds, 1-3 mm thick, possibly volcanic ash (?), at 26.7-26.75, 27.0-27.1, 27.2, and 27.25. Dark organic-rich layers appear oblique to bedding (dipping about 20°) at 26.0-26.5. <u>Spl KA2-12-27.0</u> at 27.0-27.1 (GS). <u>Spl KA2-12-27.1</u> at 27.1-27.2 (C14).
KA2-12C	27.5-27.8	Firm gray silt (catcher sample; bagged).
	27.8-28.0	Not recovered.
KA2-13	28.0-29.3	Firm gray silt with thin horizontal seams of dark gray organic-rich silt (29.0-29.3).
KA2-13C	29.3-29.6	Firm gray silt (catcher sample; bagged).
	29.6-30.5	Not recovered.
KA2-14	30.5-31.45	Very firm gray silt or clayey silt with faint lamination; no organics, except minor dark gray mottling.

KA2-14C 31.46-31.7 Very firm gray silt (catcher sample; bagged).

Maximum depth sampled 31.7 ft.

BOREHOLE KA3, PALMER HAY FLATS - 61°30'45"N, 149°30'5"W; surface elevation 19.0 ft MSL.

<u>SAMPLE</u>	<u>DEPTH (ft)</u>	<u>DESCRIPTION</u>
	0-0.3	Not recovered.
KA3-1	0.3-0.7	Frozen basal surface peat over gray silt (bagged).
	0.7-1.0	Not recovered.
KA3-2	1.0-1.25	Frozen gray silt; plugged sampler (bagged).
KA3-3	1.25-2.1	Gray to brownish-gray clayey silt. Probably disturbed at 1.25-1.7. Faint laminations at 1.7-2.1. Brownish color from minor iron staining along margins. No visible organics. <u>Spl KA3-3-2.0</u> at 1.9-2.1 (GS).
	2.1-2.8	Not recovered.
KA3-4	2.8-5.25	Wet, gray, layered and laminated clayey silt. Coarser layers (slightly darker) may contain some very fine sand. Layers are 2-15 mm thick with slight dip. Trace organic smears, otherwise no identifiable organics. Minor iron staining 5.0-5.25. Much rust on outside surface.
KA3-4C	5.25-5.5	Wet gray silt (catcher sample; bagged).
KA3-5	5.5-7.8	Very wet, gray, layered and laminated clayey silt, like KA3-4. Trace dark gray organic mottles. <u>Spl KA3-5-7.0</u> at 6.9-7.1 (GS)
	7.8-8.0	Not recovered.
KA3-6	8.0-8.65	Wet, gray, layered clayey silt like KA3-4 and 5. No organics.
	8.65-9.15	Massive wet gray clayey silt. No visible layering or organics.
KA3-7	9.15-9.5	Gray silt with no visible bedding or organics. Continuation of lower unit in KA3-6. <u>Spl KA3-7-9.3</u> at 9.2-9.4 (GS).
	9.5-11.6	Strongly laminated gray clayey silt and fine sandy silt. Organic-rich laminae at 10.05-10.15. Otherwise no organics. <u>Spl KA3-7-9.8</u> at 9.7-9.9 (GS). <u>Spl KA3-7-10.1</u> at 10.05-10.15 (C14).
KA3-7C	11.6-11.8	Gray silt (catcher sample; bagged).
	11.8-13.0	Not recovered.
KA3-8	13.0-13.6	Disturbed wet gray clayey silt and silty fine sand (probably disturbed in sampling or transport).
	13.6-15.1	Layered wet gray clayey silt and silty fine sand. No visible organics.
KA3-8C	15.1-15.4	Gray silt with one small wood fragment (catcher sample; bagged).
	15.4-15.5	Not recovered.
KA3-9	15.5-17.7	Wet, gray, silty fine sand. No visible layering except faint laminations at 17.4-17.7. Core possibly disturbed. No visible organics.
KA3-9C	17.7-18.0	Gray silty fine sand with scattered small wood fragments (catcher sample; bagged).

KA3-10	18.0-19.7	Uniform gray very fine sand. Possibly disturbed from liquefaction during sampling or transport. No organics. Channel sand? <u>Spl KA3-10-19.0</u> at 18.9-19.1 (GS).
	19.7-20.1	Not recovered.
KA3-11	20.1-22.2	Uniform gray fine sand and (at base) silty fine sand. No visible bedding except faint lamination 22.1-22.2. No visible organics.
KA3-11C	22.2-22.5	Gray fine sand (catcher sample; bagged).

Maximum depth sampled 22.5 ft.

BOREHOLE KA4, GOOSE BAY - 61°23'25"N, 149°52'10"W; surface elevation 17.1 ft MSL.

<u>SAMPLE</u>	<u>DEPTH (ft)</u>	<u>DESCRIPTION</u>
KA4-1	0-0.7	Mixed surface organics (twigs, sedge, roots) and wet gray clayey silt.
	0.7-0.8	Dark grayish brown organics (twigs, roots, sedge) with minor silt. Relatively sharp upper contact and distinctly less silt than above. Could be older peat buried by younger sediment and organics after submergence. <u>Spl KA4-1-0.7</u> at 0.7-0.8 (C14) for age of possible buried peat layer.
KA4-1C	0.8-1.2	Brown peat as above (catcher sample; bagged).
	1.2-2.9	Not recovered.
KA4-2	2.9-3.0	Slough from above - disregard.
	3.0-3.35	Brown woody peat (twigs, sedge, roots, possible sphagnum). Could be base of surface organics. Basal sample collected for date. <u>Spl KA4-2-3.3</u> at 3.25-3.35 (C14).
	3.35-4.0	Sticky gray clayey silt. No visible bedding. Minor dark gray organic mottling. <u>Spl KA4-2-3.4</u> at 3.35-3.5 (GS).
KA4-2C	4.0-4.3	Gray silt (catcher sample; bagged).
	4.3-5.5	Not recovered.
KA4-3	5.5-7.6	Firm gray silt with no visible layering and trace organics (sedge). Trace iron staining.
KA4-3C	7.6-8.0	Gray silt with trace organics (catcher sample; bagged).
KA4-4	8.0-8.75	Same as 5.5-7.6 above but no visible organics.
KA4-5	8.75-11.3	Brownish gray to gray firm clayey silt with faint lamination (9.2-9.6). Trace organics. More clay-rich at 10.1-11.3. <u>Spl KA4-5-10.0</u> at 9.9-10.1 (GS).
KA4-5C	11.3-11.5	Gray clayey silt (catcher sample; bagged).
	11.5-13.0	Not recovered.
KA4-6	13.0-14.9	Firm gray silt with minor iron staining. Numerous Desiccation cracks. No visible bedding or organics.
KA4-6C	14.9-15.2	Gray silt (catcher sample; bagged).
	15.2-15.5	Not recovered.
KA4-7	15.5-17.0	Gray to brownish gray silt with faint layering and minor iron staining. Sedge blade and twig at top, otherwise no visible organics.
	17.0-17.55	Gray to dark gray silt with organic smears and mottling. Concentrated dark gray organic zone at 17.3-17.4. Probably not a buried peat layer, but sampled for age and sedimentation rate. <u>Spl KA4-7-17.3</u> at 17.25-17.4 (C14).

KA4-7C	17.55-17.8	Gray silt (catcher sample; bagged).
	17.8-18.0	Not recovered.
KA4-8	18.0-20.15	Dry, firm gray silt with some dark gray organic-rich zones at 19.7-20.1. Breaks into hard chunks.
KA4-8C	20.15-20.4	Gray silt (catcher sample; bagged).
	20.4-20.5	Not recovered.
KA4-9	20.5-22.6	Dry, very firm, light to dark gray silt with minor organic-rich layers (charcoal?) at 21.6-21.8. Breaks into large chunks. <u>Spl KA4-9-21.8</u> at 21.7-21.85 (C14) for age and sedimentation rate (probably not a buried peat layer).
KA4-9C	22.6-22.9	Firm gray silt (catcher sample; bagged).
	22.9-23.0	Not recovered.
KA4-10	23.0-25.1	Dry, very firm, light brownish gray, gray, and some dark gray layered silt. Very brittle; breaks into chunks. Dark gray organic-rich layer at 24.15-24.25, sampled for age and sedimentation rate (not a buried peat layer). <u>Spl KA4-10-24.2</u> at 24.15-24.25 (C14).
KA4-10C	25.1-25.4	Very firm silt (catcher sample; bagged).
	25.4-25.5	Not recovered.
KA4-11	25.5-27.0	Dry, very firm and hard gray silt. Very faint layering and no visible organics.
KA4-11C	27.0-27.2	Very firm, dry silt; no organics (catcher sample; bagged).

Maximum depth sampled 27.2 ft.

BOREHOLE KA4B, GOOSE BAY - 61°23'35"N, 149°52'40"W; surface elevation 17.3 ft MSL.

<u>SAMPLE</u>	<u>DEPTH (ft)</u>	<u>DESCRIPTION</u>
KA4B-1	0-0.8	Bagged; mixed sedge blades, twigs, and water (frozen when cored) with some roots.
KA4B-1C	0.8-1.0	Frozen organic-rich silt; catcher sample; bagged (bagged).
KA4B-2C	1.0-1.4	Frozen organic-rich silt; catcher sample; bagged (bagged); Sampler plugged and pushed through soil without recovering core.
	1.4-3.0	Not recovered.
KA4B-3C	3.0-3.3	Sedge and woody peat (catcher sample; bagged). <u>Spl KA4B-3C-3.1</u> (C14).
KA4C-1C		(From adjacent borehole to attempt missed interval at 3.3-5.5 in KA4B)
	3.0-3.8	Peat; bagged.
	3.8-5.5	Not recovered.
KA4B-4	5.5-5.85	Dark brown peat (sedge, twigs, sphagnum?). Probably base of surface organics. Sampled at base for C14 age. <u>Spl KA4B-4-5.8</u> at 5.75-5.85 (C14).
	5.85-7.0	Wet gray clayey silt with no visible bedding and minor organics. <u>Spl KA4B-4-6.0</u> at 5.9-6.05 (GS).
KA4B-4C	7.0-7.3	Gray sticky clayey silt with organics (catcher sample; bagged).
KA4C-2		(From adjacent borehole to attempt missed interval in KA4B; unsuccessful)

	5.5-5.8	Same as 5.5-5.85 in KA4B-4.
	5.8-6.35	Same as 5.85-7.0 in KA4B-4.
	7.3-8.0	Not recovered.
KA4B-5	8.0-10.3	Gray clayey silt with faint layering. Trace organics at 8.25 and 9.35.
KA4C-3		(From adjacent borehole 2 ft away; same interval as KA4B-5; unsuccessful attempt to resample 7.25-8.0)
	8.0-8.5	Soft, wet organics. Probably fill-in from higher in hole (not in place).
	8.5-10.15	Wet, gray silt with minor scattered organics and some dark gray organic mottling. Twigs (1/2 in. dia.) at 9.4 and 9.9, possibly transported (although bark still intact).
	10.3-10.5	Not recovered.
KA4B-6	10.5-12.85	Moist gray clayey silt with faint layering and laminations. Void of unknown origin (not disturbed) at 10.85. Trace scattered organics. More concentrated organics (sedge?) at 12.6-12.8. Sampled for age and sedimentation rate (not a buried peat layer). <u>Spl KA4B-6-12.7</u> at 12.6-12.75 (C14).
KA4B-6C	12.85-13.2	Gray sticky silt with minor organics (catcher sample; bagged).
	13.2-13.3	Not recovered.
KA4B-7	13.3-14.25	Uniform gray clayey silt. No visible bedding; trace organic smears.
KA4B-7C	14.25-14.4	Stiff gray silt (catcher sample; bagged).
	14.4-15.5	Not recovered.
KA4B-8	15.5-17.1	Uniform gray clayey silt. Minor crudely layered but distorted dark gray organic-rich layers at 16.9-17.0.
KA4B-8C	17.1-17.3	Gray silt (catcher sample; bagged).
	17.3-18.0	Not recovered.
KA4B-9	18.0-20.5	Gray to dark gray clayey silt. Several areas of dark gray organic-rich mottling and layering (18.2-18.3, 18.9-19.0, 19.2-19.3, 19.4).
KA4B-10	20.5-22.9	Very firm, nearly dry, gray to light brownish gray clayey silt. Distinct but disturbed layering. Light brownish gray zones along cracks and partings, possibly from iron staining. No visible organics, but darker gray zones may be slightly organic-rich. Concooidally fractured.
KA4B-11	22.9-25.4	Same as KA4B-10. Several small (up to 1/4 in.) subrounded pebbles at 24.8. No visible organics. <u>Spl KA4B-11-24.8</u> at 24.7-24.9 (GS).

Maximum depth sampled 25.4 ft.

BOREHOLE KA5, PALMER HAY FLATS - 61°31'40"N, 149°26'55"W; surface elevation 21.4 ft MSL.

<u>SAMPLE</u>	<u>DEPTH (ft)</u>	<u>DESCRIPTION</u>
	0-4.7	Not sampled; organics and water.
KA5-1	4.7-5.1	Brown to dark brown woody, sedge, and sphagnum(?) peat. Sharp base. <u>Spl KA5-1-5.0</u> at 5.0-5.1 (C14).
	5.1-6.0	Wet, gray, layered silt with trace organics. <u>Spl KA5-1-5.6</u> at 5.5-5.7 (GS).
	6.0-6.25	Wet, gray, clayey silt with faint layering and no visible organics. <u>Spl KA5-1-6.2</u> at 6.1-6.25 (GS).

KA5-1C	6.25-6.5	Gray silt with minor organics (catcher sample; bagged).
KA5-2	6.5-8.2	Wet, gray, laminated silt and clayey silt. Laminae dip 10-30° at 6.5-7.3. No visible organics.
KA5-2C	8.2-8.6	Gray silt (catcher sample; bagged).
KA5-3	8.6-8.95	Wet, gray, laminated silt and clayey silt. No visible organics.
KA5-4	9.0-9.7	Wet, gray silt and clayey silt with distorted laminae.
	9.7-11.4	Wet, gray, laminated and thin-layered silt and clayey silt. Dark gray silt layer at 10.25-10.35 may be slightly organic-rich. Dark gray organic-rich(?) mottling at 10.6-11.0. <u>Spl KA5-4-10.0</u> at 9.9-10.1 (GS).
KA5-4C	11.4-11.8	Gray silt (catcher sample; bagged).
KA5-5	11.8-13.2	Wet, gray, laminated and thin-layered silt and clayey silt. Disturbed at 11.8-12.7. No visible organics.
KA5-6	13.2-15.7	Moist, gray, laminated and thin-layered silt and clayey silt. No visible organics. Cross-bedded fine-sandy layers at 14.5-14.7. <u>Spl KA5-6-15.3</u> at 15.2-15.4 (GS), silt. <u>Spl KA5-6-15.6</u> at 15.55-15.7 (GS), clayey silt.
KA5-6C	15.7-16.0	Gray silt (catcher sample; bagged).
KA5-7	16.0-16.5	Gray silt (bagged).
KA5-8	16.5-18.7	Moist, gray, layered silt and clayey silt. Layering less distinct than in KA5-6. No visible organics.
	18.7-18.9	Uniform gray clay or clayey silt. No visible layering or organics.
KA5-8C	18.9-19.0	Gray silt (catcher sample; bagged).
KA5-9	19.0-19.95	Uniform gray clay or clayey silt. No visible layering or organics. Continuation of layer starting at 18.7.
KA5-10	19.95-22.45	Firm, uniform gray silty clay. No visible bedding, except faint dark thin (< 1/8 in.) gray organic-rich(?) layer at 21.6. <u>Spl KA5-10-21.0</u> at 20.9-21.1 (GS).
KA5-10C	22.45-22.7	Gray silt (catcher sample; bagged).
	22.7-24.0	Not recovered.
KA5-11	24.0-25.95	Same as KA5-10. Slightly organic-rich (dark gray) layer at 25.4-25.5. Small piece of twig at 25.4.
KA5-11C	25.95-26.2	Gray silty clay (catcher sample; bagged).
	26.2-26.5	Not recovered.
KA5-12	26.5-27.0	Same as KA5-10 and KA5-11.
	27.0-29.0	Not recovered. Sample slid out of tube during recovery.
KA5-13	29.0-30.85	Gray silty clay, same as KA5-10, 11, and 12. Very faint darker gray zones, possibly organic-rich (for example, 30.1-30.15). <u>Spl KA5-13-30.0</u> at 29.9-30.1 (GS).
KA5-13C	30.86-31.1	No change (catcher sample; bagged).

Maximum depth sampled 31.1 ft.

BOREHOLE KA6, PALMER HAY FLATS - 61°31'20"N, 149°15'25"W; surface elevation 18.7 ft MSL.

<u>SAMPLE</u>	<u>DEPTH (ft)</u>	<u>DESCRIPTION</u>
	0-0.5	Ice.
	0.5-1.9	Frozen silt and organics (not recovered).
KA6-1	1.9-2.2	Base of frozen silt (bagged).
KA6-2	2.2-2.35	Sedge and woody peat mixed with some gray silt. Could be basal surface peat, base of older peat, or fill-in from borehole wall above. <u>Spl KA6-2-2.3</u> at 2.2-2.35 (C14).
	2.35-2.95	Uniform gray silty clay with scattered organics. Slightly organic-rich layer at 2.6-2.7. Sharp base. <u>Spl KA6-2-2.9</u> at 2.8-2.95 (GS).
	2.95-3.2	Brown to dark brown sedge and sphagnum(?) peat. Sharp upper and lower contacts. <u>Spl KA6-2-3.0</u> at 2.95-3.05 (C14).
	3.2-3.4	Uniform gray clayey silt. No organics.
KA6-2C	3.4-3.7	Gray clayey silt (catcher sample; bagged).
	3.7-4.0	Not recovered.
KA6-3	4.0-4.3	Gray clayey silt with scattered organics and very faint layering.
	4.3-4.375	Very dark brown compressed sedge or sphagnum peat. Sharp top and bottom. <u>Spl KA6-3-4.35</u> at 4.3-4.375 (C14).
	4.375-4.45	Light brownish gray organic-rich silt or ash(?). <u>Spl KA6-3-4.4</u> for microscopy.
	4.45-4.55	Dark brown compressed sedge or sphagnum peat. Could be slightly older submerged layer (2 events?). Sharp base, gradational top.
	4.55-5.15	Gray clayey silt with very faint layering and scattered organics.
KA6-3C	5.15-5.4	Gray silt (catcher sample; bagged).
	5.4-6.5	Not recovered.
KA6-4	6.5-7.3	Uniform gray clayey silt with no visible bedding or organics.
	7.3-7.4	Laminated gray clayey silt and silty fine sand. Minor dark gray organic(?) smear.
	7.4-7.525	Gray clayey silt with no visible bedding. Minor gray organic(?) smear.
	7.525-7.575	Grayish brown organic-rich silt. <u>Spl KA6-4-7.55</u> at 7.525-7.575 (C14).
	7.575-7.8	Gray clayey silt with scattered organics. No visible bedding.
	7.8-7.9	Dark brown compressed peat. Sharp top and bottom. <u>Spl KA6-4-7.85</u> at 7.8-7.9 (C14).
KA6-4C	7.9-8.1	Gray silt (catcher sample; bagged).
	8.1-9.0	Not recovered.
KA6-5	9.0-9.8	Very wet, disturbed clayey silt with faint layering and minor dark gray organic smears.
KA6-6	9.8-10.7	Same as KA6-5.
	10.7-11.5	Not recovered.

KA6-7	11.5-13.85	Wet gray clayey silt with faint disturbed bedding and scattered organics. Pocket of sedge(?) at 13.1. Mottled with dark gray smears, possibly fine organics. <u>Spl KA6-7-12.0</u> at 11.0-12.1 (GS).
	13.85-14.0	Not recovered.
KA6-8	11.5-7	Extra material from unknown source (structure precludes fill-in from borehole walls). Recovered at same apparent interval as KA6-7. Wet gray clayey silt with scattered organics. Some distinct layering in lower half. Upper part disturbed.
KA6-9	14.0-16.45	Wet, uniform gray clayey silt with trace scattered organics and slight iron staining at 16.2-16.45.
KA6-10	fill-in	Gray clayey silt with some organics.
KA6-11	16.5-19.0	Gray clayey silt with faint layering and some iron staining at 16.5-18.4. Distinct layering and lamination at 18.4-18.95, and no iron staining. Trace scattered organics.
KA6-12	19.0-20.95	Gray silty clay with very faint layering and no visible organics. Several thin (1/8 in.) fine-sandy partings at 20.2-20.3. <u>Spl KA6-12-20.0</u> at 19.9-20.1 (GS).
	20.95-21.5	Not recovered.
KA6-13	21.5-23.15	Firm, gray clayey silt with very faint layering and no visible organics.
KA6-13C	23.15-23.4	Gray clayey silt (catcher sample; bagged).
	23.4-24.0	Not recovered.
KA6-14	24.0-25.65	Firm, gray clayey silt. Very faint layering and no visible organics at 24.0-25.2. Distinct layering accentuated by dark gray organic smears at 25.2-25.65. Darkest layer at 25.25-25.3. <u>Spl KA6-14-25.3</u> at 25.25-25.35 (C14).
KA6-14C	25.65-25.9	Gray clayey silt (catcher sample; bagged).
	25.9-26.5	Not recovered.
KA6-15	26.5-28.75	Firm, gray to dark gray clayey silt with no visible bedding. Numerous dark gray organic smears.
KA6-15C	28.75-29.0	Firm gray clayey silt with 1/2 in. dark gray organic-rich layer.
KA6-16	29.0-31.5	Firm, gray clayey silt with minor dark gray organic smears. No visible bedding. <u>Spl KA6-16-30.0</u> at 29.9-30.1 (GS).
KA6-17	31.5-33.75	Same as above. Thin (1/4 in.) dark gray organic-rich(?) layer at 32.75. Trace fine sandy lenses at (e.g. 31.7, 32.65).
KA6-17C	33.75-34.0	Gray clayey silt with faint dark gray organic smears near top (catcher sample; bagged).
KA6-18	34.0-36.5	Firm gray clayey silt with minor dark gray organic smears. Numerous light gray fine sandy lenses and partings at 35.9-36.5.
KA6-19	36.5-38.2	Firm, gray clayey silt with trace dark gray organic smears, distorted faint layering, and several light gray fine-sandy lenses.
	38.2-38.55	Interlayered grayish brown silty fine sand and brownish gray to gray fine-sandy silt and clayey silt. Slightly distorted layering. No visible organics. <u>Spl KA6-19-38.4</u> at 38.35-38.45 (GS).
	38.55-39.0	Uniform gray silty clay with no visible bedding or organics. Trace fine sand. <u>Spl KA6-19-38.8</u> at 38.7-38.9 (GS).

KA6-20	39.0-39.6	Gray silty clay with faint distorted layering; probably disturbed. No organics.
	39.6-40.2	Gray laminated clayey silt and silty fine sand. Laminae tilted and distorted at 39.6-39.9. No visible organics.
	40.2-40.45	Gray to grayish brown silty fine sand. No visible bedding or organics.
	40.45-41.25	Soft gray silt and clayey silt with distorted bedding. No visible organics.
KA6-21	41.25-41.9	Soft gray clayey silt with interbeds of silty fine sand. Distorted bedding at 41.25-41.6. Less disturbed at 41.6-41.9. Layering is horizontal at base. Possibly trace organics at 41.85.
	41.9-42.35	Grayish brown fine sand with distorted interbeds of gray clayey silt. No visible bedding.
	42.35-43.35	Distorted interbeds of gray clayey silt and silty fine sand. No visible organics.
	43.55-43.75	Horizontal interbeds of gray fine sand and clayey silt. No visible organics.
KA6-21C	43.75-44.0	Firm gray silty fine sand (catcher sample; bagged).

Maximum depth sampled 44.0 ft.


RIVER BLUFF SECTION KA7, PALMER HAY FLATS - loc. near KA1; surface elev. about 18 ft MSL.

<u>SAMPLE</u>	<u>DEPTH (ft)</u>	<u>DESCRIPTION</u>
	0-3.0	Not sampled; grayish brown silt with plant fragments and roots near surface.
KA7-3.0	3.0	Flattened wood pieces and twigs lying horizontally in a discontinuous layer up to 1 in. (2.5 cm) thick. <u>Spl KA7-3.0</u> at 3.0 ft (C14).
KA7-3.5	3.5	Sedge and woody peat in a laterally extensive but discontinuous layer up to 1 in. (2.5 cm) thick. <u>Spl KA7-3.5</u> at 3.5 ft (C14).

APPENDIX C
GRAIN-SIZE ANALYSES


Sample-Depth(ft)	Mean Dia.(mm)	Mean Dia.(ϕ)*	Standard Deviation*	Skewness*	Kurtosis*	Percent Gravel	Percent Sand	Percent Silt	Percent Clay
TA1-17.1	0.0122	6.36	1.74	1.51	4.43	0.00	0.00	83.73	16.27
TA1-19.0	1.5801	-0.66	1.56	0.79	3.56	61.71	36.71	1.58	0.00
TA1-25.3	0.0043	7.85	1.79	0.41	2.31	0.00	0.00	58.21	41.79
TA1-26.3	0.0047	7.74	1.87	0.28	2.43	0.00	0.90	58.50	40.60
TA1-33.0	0.0060	7.38	2.11	0.51	2.13	0.00	0.40	64.00	35.60
TA1-36.6	0.0126	6.31	2.14	1.09	3.08	0.00	5.00	74.00	21.00
TA1-41.8	0.0166	5.91	1.94	1.23	3.29	0.00	4.00	78.70	17.30
TA1-45.6	0.0209	5.58	1.68	1.66	4.89	0.00	4.70	83.70	11.60
TA1-57.0	0.0337	4.89	1.27	2.45	8.77	0.00	11.80	82.40	5.80
TA1-61.0	0.0270	5.21	1.30	2.42	9.07	0.00	4.20	89.60	6.20
TA8-4.7	0.1321	2.92	0.79	0.16	2.33	0.00	85.94	14.06	0.00
TA8-9.5	0.1058	3.24	0.57	0.38	2.47	0.00	85.43	14.57	0.00
TA8-12.5	0.0145	6.11	1.65	1.19	3.59	0.00	2.00	83.10	14.90
TA8-16.7	0.0126	6.31	1.45	1.38	4.18	0.00	0.00	85.94	14.06
TA8-20.0	0.0154	6.02	1.45	1.87	5.17	0.00	0.00	88.81	11.19
TA8-24.6	0.0212	5.58	1.18	2.04	6.92	0.00	1.10	91.80	7.10
TA8-30.2	0.0182	5.78	1.32	1.50	4.60	0.00	1.20	89.00	9.80
TA8-35.0	0.0137	6.19	1.72	1.32	3.65	0.00	0.20	83.50	16.30
TA8-45.5	0.0050	7.64	1.92	0.43	2.20	0.00	0.20	61.30	38.50
TA8-50.5	0.0047	7.73	1.62	0.43	2.59	0.00	0.00	61.53	38.47
KA1B-3.1	0.0043	7.87	2.03	0.34	2.04	0.00	0.00	56.33	43.67
KA1-5.7	0.0044	7.82	2.03	0.56	2.44	0.00	0.10	59.30	40.60
KA1-10.0	0.0056	7.48	1.70	0.35	2.03	0.00	0.00	63.96	36.04
KA1-11.6	0.0052	7.59	2.07	0.64	2.47	0.00	0.00	63.60	36.40
KA1-17.9	0.0067	7.23	1.86	0.83	2.74	0.00	0.00	71.20	28.80
KA1-25.0	0.0039	7.99	2.05	0.39	2.22	0.00	0.20	54.90	44.90
KA2-3.5	0.0140	6.16	1.50	1.43	4.47	0.00	0.80	86.30	12.90
KA2-4.0	0.0042	7.91	1.97	0.24	2.10	0.00	0.90	53.70	45.40
KA2-7.0	0.0148	6.08	1.61	1.54	4.51	0.00	0.40	86.00	13.60
KA2-20.0	0.0034	8.18	1.96	0.33	2.26	0.00	0.00	50.60	49.40
KA2-27.0	0.0040	7.95	2.26	0.40	2.11	0.00	0.30	55.20	44.50
KA3-2.0	0.0091	6.78	1.83	1.19	3.37	0.00	0.00	78.00	22.00
KA3-7.0	0.0067	7.23	2.21	1.02	2.92	0.00	0.00	71.10	28.90
KA3-9.3	0.0045	7.80	1.94	0.42	2.06	0.00	0.00	58.29	41.71
KA3-9.8	0.0158	5.98	1.57	2.12	6.96	0.00	0.00	89.41	10.59
KA3-19.0	0.0652	3.94	0.88	1.62	0.69	0.00	56.40	42.90	0.70
KA4-3.4	0.0039	7.99	1.83	0.24	2.20	0.00	0.00	53.20	46.80
KA4-10.0	0.0057	7.46	1.73	0.54	2.57	0.00	0.20	66.20	33.60
KA4B-6.0	0.0036	8.12	2.06	0.48	2.27	0.00	0.00	54.00	46.00
KA4B-24.	0.0044	7.83	2.13	0.48	2.25	0.00	0.00	58.00	42.00
KA5-5.6	0.0119	6.39	1.68	1.15	3.37	0.00	1.00	81.80	17.20
KA5-6.2	0.0056	7.49	1.70	0.45	2.53	0.00	0.00	64.70	35.30
KA5-10.0	0.0075	7.06	1.71	0.96	3.02	0.00	0.00	74.48	25.52
KA5-15.3	0.0157	5.99	1.71	1.96	6.12	0.00	0.00	88.02	11.98
KA5-15.6	0.0036	8.13	2.01	0.25	2.26	0.00	0.10	50.30	49.60
KA5-21.0	0.0029	8.45	1.79	0.21	2.52	0.00	0.10	42.70	57.20
KA5-30.0	0.0023	8.78	1.88	0.02	2.45	0.00	0.20	35.40	64.40
KA6-2.9	0.0026	8.59	1.77	-0.19	2.51	0.00	0.60	36.50	62.90
KA6-12.0	0.0076	7.04	1.94	0.93	2.89	0.00	0.50	74.10	25.40
KA6-20.0	0.0022	8.80	1.88	0.21	2.26	0.00	0.00	37.91	62.09
KA6-30.0	0.0045	7.79	1.87	0.31	2.23	0.00	0.00	57.80	42.20
KA6-38.4	0.0203	5.62	1.94	1.59	4.51	0.00	6.40	79.40	14.20
KA6-38.8	0.0023	8.79	2.12	-0.25	2.20	0.00	0.00	34.00	66.00


* $\Phi(\phi) = -\log_2(d_{mm})$. Standard deviation, skewness, and kurtosis are in phi units.





KEY:

- Sand and gravel.
- Silt or fine sand with negligible clay.
- Silt and clay.
- Peat.
- Organic-rich sediment (contains plant fragments or dark gray, disseminated fine organic matter).
- No sample recovery.
- Radiocarbon age (y: BPI) and one-sigma range, uncalibrated.
- Approximate mean higher high water.
- Mean grain diameter and one standard deviation.







- KEY:
- Silt or fine sand with negligible clay.
 - Silt and clay.
 - Peat.
 - Organic-rich sediment (contains plant fragments or dark gray, disseminated fine organic matter).
 - No sample recovery.
 - Approximate mean higher high water.
 - Mean grain diameter and one standard deviation.

