ALASKA DIVISION OF GEOLOGICAL & GEOPHYSICAL SURVEYS
IN COOPERATION WITH U.S. BUREAU OF MINES

PUBLIC-DATA FILE 94-55
PLATE 1

GENERALIZED GEOLOGIC MAP OF THE WESTERN ENDICOTT MOUNTAINS, CENTRAL BROOKS RANGE, ALASKA

By
C.G. Mull and M.B. Werdon

1994

DESCRIPTION OF MAP UNITS

SURFICIAL DEPOSITS
Modified from Hamilton, 1980

Qal ACTIVE ALLUVIAL DEPOSITS (Quaternary)–Silt, sand and gravel deposited by modern streams; includes flood-plain deposits which are lightly vegetated in some areas.

Qaf ALLUVIAL FAN DEPOSITS (Quaternary)–Poorly to moderately sorted fluvial silt, sand and gravel forming dotted fans and aprons where tributaries join higher order streams.

Qu SURFICIAL DEPOSITS, UNDIVIDED (Quaternary)–Includes, colluvium and older alluvium; alluvial deposits above present stream levels, alluvial fans, and upland silt deposits.

Qm MORAINAL DEPOSITS (Quaternary)–Poorly sorted unstratified till ranging from clayey stony silt to sandy boulder gravel; contains large erratic boulders in some places; deposits are commonly hummocky, may be heavily vegetated, locally forms terminal moraines but in some places form well defined ridges and concentric ridges.

Qgu GLACIAL DEPOSITS, UNDIFFERENTIATED (Quaternary)–Stratified and unstratified drift of various glacial intervals. Includes till, outwash and ice-contact deposits.

QTt TERTIARY GRAVEL (Quaternary and Tertiary)–Gravel and sandy gravel, forms terraces ranging from 3-4 m above modern stream levels, to high-level terraces.

FOREDEEP DEPOSITS OF THE COLVILLE BASIN

Kto TOROK SHALE (Lower Cretaceous, Albian)–Soft dark-gray to black clay shale and thinly interbedded silty shale, generally exposed only in discontinuous stream cutbanks, most exposures lightly baked, lower portion interfingers with Fortress Mountain Formation.

Kfm FORTRESS MOUNTAIN FORMATION, UNDIVIDED (Lower Cretaceous, Albian)–Dominantly greenish-gray graywacke interbedded with dark-gray mudstone and shale, interfingers with Torok Shale, in places contains differentiated units:

Kfcg–As above, contains thick massive beds of pebble- to boulder- conglomerate, clasts are well rounded and consist dominantly of gray, black, and green chert and a large variety of altered mafic igneous rocks.

ALLOCHTHONOUS ROCKS

ENDICOTT MOUNTAINS ALLOCHTHON

KEY CREEK, AND IVOTUK HILLS SEQUENCES UNDIVIDED

Named for exposures in the western Delong Mountains (Mayfield and others, 1968) and for exposures in the Killik River quadrangle Mull and others, 1994). In map area, exposed along mountain front in eastern part of area, in Ivotuk Hills area, and in structural windows at west end of map area at Safari Creek and at head of tributary of Anuk River.

KMu CRETACEOUS TO MISSISSIPPIAN ROCKS, UNDIVIDED—mapped in structural window at headwaters of north fork of Anuk River in western part of map area, and north of mountain near Story Creek,. Generally intensely deformed thrust slivers and isolated exotic blocks of Mississippian to Triassic rocks in area too small to be delineated at scale of map, in some places probably in a groundmass of Cretaceous mudstone and graywacke.

Ko CHOPKIRUAK FORMATION (Cretaceous, Neocomian)–dark gray mudstone, graywacke, and local cobble conglomerate; generally intensely deformed, thickness unknown.

Kcg LIMESTONE CONGLOMERATE (Lower Cretaceous Neocomian)–Gray to reddish gray, thin-bedded limestone, interbedded with dark gray to black shale. Limestone is 1-2 m thick, composed entirely of the pelecypod *Buchia sublaevis* of Valanginian age, commonly weathers a distinctive reddish brown color. Limestone is commonly highly compacted and fossils unrecognizable, unit is identified by distinctive wavy laminated appearance on surface broken across bedding.

JPe ETIVLUK GROUP (Middle Jurassic to Permian)–Divided into:

JTRa OTUK FORMATION (Middle Jurassic to Triassic)–Interbedded fossiliferous black chert, shale, and thin bedded black and tan banded limestone characterized by abundant *Monotis* ss. or *Halobia* sp. pelecypods. The Jurassic Blankenship Member at the top is locally exposed and consists of organic-rich black shale and thin bedded chert. Formation is commonly exposed only as yellow-gray weathering limestone rubble or in structurally complicated partial sections in stream cuts. Thickness <100 m.

Ps SIKSIKPUK FORMATION (Permian)–Dominantly nonresistant thin bedded siltstone, shale, and minor silicified mudstone. Consists of a distinctive thin yellowish-orange weathering basal greenish-gray siltstone that grades up into maroon- to light-green mottled siltstone with scattered thin barite seams and crystal aggregates, a resistant gray chert-pebble sandstone member, and locally a <10 cm thick, and poorly exposed gray argillite. Formation is commonly exposed only as yellow-orange weathering rubble on hill sides or in stream cuts where it is recognized by the conspicuous yellowish-orange weathering beds that overlie Lisburne Group. Thickness <100 m.

Mld LISBURNE GROUP, DOLOMITE FACIES (Upper Mississippian) (Dumoulin and Harris, 1994)–Dolomitic, fine- to medium-grained, thick-bedded, massive, with white to bluish-gray weathering chert, some bioclastic beds, some beds of upsection mudstone and chert, dominantly thin-bedded chert and minor shale in lower part. In some zones, contains abundant pinpoint to vuggular porosity filled with tarry oil and solid hydrocarbons. Forms high rubble covered ridges of Ivotuk Hills 20 km north of mountain front in northwestern part of map area. Thickness about 225 m (Dumoulin and Harris, 1994).

Mkn KUNA FORMATION, OF LISBURNE GROUP (Lower and Upper? Mississippian)–Interbedded black sooty siliceous mudstone, and shale, minor limestone and dolomite; also contains light-gray bioclastic limestone in western part of map area. Unit is generally shaly and poorly exposed, forms low rubble covered slopes, in some areas where limestone is present forms low resistant ridges near mountain front in eastern part of map area. Type section of formation is located in Sallat Creek tributary of Kuna River in western part of map area; reference section for formation is located on Chik Creek 2.5 km north of mountain front in eastern part of map area. Contains Lower Mississippian (Osagean) conodonts in Howard Pass quadrangle (Dumoulin and others, 1994). Thickness <75 m. In places bioclastic limestone is undifferentiated:

Mkns KUNA FORMATION, LIMESTONE FACIES (Lower Mississippian)–Light-gray weathering bioclastic limestone, appears to interfinger with sooty black shale in the Kuna Formation. Limestone is commonly thin bedded and forms ridges 1-2 km in front, forms scattered generally low rubble-covered slopes. In many areas includes poorly exposed Kayak Shale at base. Contains Lower Mississippian (Kinderhookian) conodonts (Anita Harris, written communication, 1994). Thickness <20 m.

KAYAK SHALE (Lower Mississippian)–Dark-gray to black fissile clay shale with conspicuous yellowish-brown-weathering thin fossiliferous limestone beds near top, commonly contains conspicuous reddish-brown weathering nodules; lower part contains interval characterized by greenish-gray siltstone and thinly interbedded phyllitic shale; interval becomes thicker and more prominent west of Etivluk River, both interbedded with head of north fork of Anuk River, contains numerous thin siltstone beds interbedded with shale, and a unit of thin bedded black chert and shale near base of formation. West of Etivluk River, unit contains extrusive volcaniclastic and apparently intrusive felsic rocks that become more abundant westward. Unit is generally poorly exposed on bare black-weathering hillsides or in stream cuts where it is completely deformed by basinal folding. Contains Lower Mississippian (Kinderhookian) conodonts in Howard Passs quadrangle (Dumoulin, 1980; Dumoulin and others, 1994). Igneous rocks locally differentiated as:

Mv VOLCANIC ROCKS (Lower Mississippian)–Extrusive volcaniclastic rocks, felsic to intermediate; range from coarse agglomerate to rhyolitic tuff, light gray to dark-green weathering. Unit is best exposed in Kayak Shale sequence in Ivotuk Hills, associated with Kayak Shale in a number of localities in Fay Creek and Tuluto Creek area north of mountain front, and in structural window at head of tributary of Anuk River. Gradationally overlain by black phyllitic Kayak Shale at one locality on upper Tuluto Creek. Thickness probably >100 m in some areas.

MDk KANAYUT CONGLOMERATE (Lower Mississippian? to Upper Devonian)–Thick succession of gray- to brown-weathering quartzitic sandstone and minor quartz- and chert-pebble conglomerate, forms highest mountain peaks in western Endicott Mountains. At eastern edge of map area north of Toyuk thrust unit is probably >900 m thick, thins rapidly to north and west of East Fork Etivluk River; south of Toyuk thrust in southeast corner of map is 228 m thick (J.T. Moore, written communication, 1994). North of Toyuk thrust is differentiated into:

Dkal SHAININ LAKE MEMBER (Upper Devonian)–Quartzitic sandstone and minor pebble conglomerate, weathers rusty forms steep dark gray to black weathering mountain ridges, probably includes unrecognized Stuver Member at top, grades down into Ear Peak Member.

Dke EAR PEAK MEMBER (Upper Devonian)–Shale, siltstone, quartzitic sandstone, and conglomerate, forms steep brown- to gray-weathering slopes with resistant ridges and cliffs.

Dn NOATAK SANDSTONE (Upper Devonian)–Light gray- to light-brown, fine- to coarse-grained thick bedded sandstone and quartzite in units up to 20 m thick; interbedded with light-gray to brown shale with ironstone nodules; sandstone commonly crossbedded, contains scattered limonitic spots, upper part generally forms steep ledge-weathering rubble-covered slopes, lower part generally bedded downward in section. South of Toyuk thrust formation is locally differentiated into:

Dnd NOATAK SANDSTONE, LOWER PART (Upper Devonian)–Gray to greenish-gray, medium-bedded sandstone, generally fine-grained, calcareous, finely micaceous, contains abundant sedimentary structures including crossbedding, micaceous laminations, and ripple marks; beds up to 2 m thick, interbedded conspicuous micaceous shale and local thin silty limestone beds; interval forms generally smooth, dark reddish-brown to brown weathering, rubble-covered slopes in mountains northeast of Ngo River , probably >500 m thick. At base contains Late Devonian (Famennian) brachiopods and ammonoids (J. T. Dutro and others, in press).

Dnc CONGLOMERATE AND SANDSTONE UNIT, UNNAMED (Upper Devonian)–Massive, thick-bedded, white to light-gray weathering pebble conglomerate and quartzitic sandstone; conglomerate contains matrix-supported white quartz and black and gray chert pebbles to 2 cm diameter; sandstone is dominantly fine- to medium-grained, contains abundant sedimentary structures, and locally contains abundant yellowish-brown limonite spots in places. Local conspicuous cliff-forming marker horizon up to 25 m thick between Ivotuk Creek and Iteriak Creek area northeast of Kilik River Quadrangle.

Dnh NOATAK SANDSTONE AND HUNT FORK SHALE, UNDIVIDED (Upper Devonian)–Transitional facies south of Toyuk thrust in south-western part of Killik River Quadrangle; consists of gray to greenish-gray medium-bedded micaceous and calcareous sandstone as in overlying Noatak Sandstone, beds up to 1 m thick, interbedded with dominantly silty and gray silty micaceous shale; interval forms generally smooth dark reddish-brown to brown weathering mountain slopes with local micaceous sheen. Thickness probably about 300 m.

Dhf HUNT FORK SHALE (Upper Devonian)–Dark gray- and olive-green shale, with interbedded thin bedded quartz-chert wacke, quartzite, sandstone, and minor conglomerate in upper part north of Toyuk thrust; south of Toyuk thrust, unit grades progressively downward into a thick lower part. South of Toyuk thrust, unit grades progressively downward into a thick lower part. South of Toyuk thrust, unit grades progressively downward into interbedded greenish-gray micaceous sandstone and phyllite into dominantly micaceous phyllite that forms smoothly weathered slopes with a conspicuous micaceous sheen, forms base of exposed sequence. Thickness > 500m.

ANIUK RIVER SEQUENCE

Here named for the Aniuk River area on the south side of the western Endicott Mountains in western part of map area.

Ko CHOPKIRUAK FORMATION (Lower Cretaceous)–Dark gray shaly mudstone and thin-bedded fine grained graywacke, at base locally contains thin interval of maroon and greenish gray clay shale that may be correlative with Valanginian ecquonid limestone horizon on Key Creek and Ivotuk Hills sequences, but no limestone observed. Exposed only locally in rubble-covered slopes on north side of Rough Mountain Creek north of Ishkkuk Mountain, and in stream cutbank southwest of Ialuit Mountain.

JPe ETIVLUK GROUP (Middle Jurassic to Permian)–exposed in bare slopes and stream cuts in broad valleys east and west of Nigtun Lake in Howard Pass quadrangle, subdivided locally differentiated into:

JTRa OTUK FORMATION (Middle Jurassic to Triassic)—Thin-bedded black chert, shale, and thin-bedded black and tan banded limestone characterized by abundant *Monotis* ss. or *Halobia* sp. pelecypods. The Jurassic Blankenship Member at the top of the formation has not been recognized on this sequence owing to poor exposure but is probably present. Exposed in bare slopes east of Nigtun Lake, in valley southwest of Ialuit Mountain, and north of Rough Mountain Creek.

Ps SIKSIKPUK FORMATION (Permian)–Chert and siliceous mudstone, green to greenish gray, sreaky bedded in 2-10 cm thick beds, underlain by maroon to greenish-gray siliceous and phyllitic shale, unit <30 m thick. Exposed locally on north side of Rough Mountain Creek north of Ishkkuk Mountain, and in bare slopes in area of headwater tributaries of Fanta Creek.

Mkn KUNA FORMATION (Lower Mississippian)–soft black sooty shale and minor silicified mudstone, generally poorly exposed, <10 m thick, contains lower Mississippian (Osagean) conodonts (Dumoulin and Harris, 1994). Light gray bioclastic limestone at base locally differentiated as:

Mlmd BASAL LIMESTONE OF KUNA FORMATION (Lower Mississippian)–Bioclastic limestone, light gray, fine to medium grained, abundant crinoidal debris, in beds 10 cm to 1 m thick, forms ledges 1-2 m high in bare slopes. Unit is <10 m thick, in part contains gray to greenish-gray silty limestone, weathers yellowish-brown. Disconformously exposed as low rubble covered ridges and hilltops north of Rough Mountain Creek, on Mountain of Ialuit Mountain, and along south side of mountains between Ipnavik River and Safari Creek. Contains Lower Mississippian (Osagean) conodonts (Anita Harris, written communication, 1993). Unit <10 m thick.

Mi ISIKUT UNIT (Lower Mississippian)–dominantly greenish-gray thin bedded, platy siltstone and interbedded phyllitic shale; siltstone commonly flaser bedded, contains abundant sedimentary structures including crossbedding, micaceous laminations, and ripple marks. Unit is marked by light-gray weathering quartzose sandstone member in Black phyllitic shale. Overlying beds contain scattered very fine-grained sandy limestone beds to 2 m thick that commonly contain bioturbated and brecciated crinoidal debris. Upper part of unit contains increasing amounts of phyllitic shale. Locally contains minor undifferentiated felsic intrusive rocks. Unit is overlain with sharp contact by Kanayut Conglomerate and Noatak Sandstone Creek Sequence. Contains basal Mississippian (Kinderhookian) conodonts and Mississippian (Osagean) conodonts (Anita Harris, written communication, 1994; John Carter, written communication, 1994). Forms conspicuous reddish-brown to orange-brown weathering bare rubble-covered slopes along south side of western Endicott Mountains from Isikut Mountain west to the north fork of Anuk River, and on Picnic Creek allochthon on mountains from Story Creek to Safari Creek. Measured thickness 450 m on north side of Isikut Mountain (68°12.7'N, 157°41.5'W); 350 m on Mountain of Ialuit (68°12.7'N, 157°41.5'W) (T.E. Moore, written communication, 1994). Areas with more abundant igneous rocks locally differentiated as:

Miv ISIKUT UNIT WITH VOLCANIC ROCKS UNDIVIDED (Mississippian)–Isikut unit as above, but with interbedded light-gray to greenish gray extrusive volcaniclastic rock and dark green possibly intrusive, mafic igneous rock; mapped on north side of mountains between Ipnavik River and Safari Creek and locally on south side of mountains.

MDkn KANAYUT CONGLOMERATE AND NOATAK SANDSTONE, UNDIVIDED—greenish-gray quartzitic sandstone, calcareous sandstone, and minor conglomerate. In Anuk River Sequence, Kanayut Conglomerate and Noatak Sandstone appear to be lithologically similar to the units in the Killik River and Key Creek Sequences in the eastern Endicott Mountains, but are much thinner. Forms high mountains between Ipnavik River and Fenian Lake. In northern part of mountains, imbricated by numerous thrust faults and difficult to differentiate owing to rubble cover; structurally less complicated in trend south of upper Ipnavik River near southern edge of map area. Locally differentiated into:

MDk KANAYUT CONGLOMERATE (Lower Mississippian? to Upper Devonian)–Dark gray to black, medium- to thick-bedded quartzitic sandstone and quartz-pebble conglomerate; local green- pebble conglomerate, supports heavy growth of black lichens, interfingers downward with Noatak Sandstone. In western part of map area northeast of Fenian Lake and upper Story Creek area, top of unit consists of about 75 m of several fine-grained quartzitic sandstone intervals to 5 m thick, that form a prominent marker horizon. Unit is generally poorly exposed on bare black-weathering hillsides or in stream cuts where it is completely deformed by basinal folding. Contains Lower Mississippian (Kinderhookian) conodonts in Howard Passs quadrangle (Dumoulin, 1980; Dumoulin and others, 1994). Igneous rocks locally differentiated as:

Mv VOLCANIC ROCKS (Lower Mississippian)–Extrusive volcaniclastic rocks, felsic to intermediate; range from coarse agglomerate to rhyolitic tuff, light gray to dark-green weathering. Unit is best exposed in Kayak Shale sequence in Ivotuk Hills, associated with Kayak Shale in a number of localities in Fay Creek and Tuluto Creek area north of mountain front, and in structural window at head of tributary of Aniuk River. Gradationally overlain by black phyllitic Kayak Shale at one locality on upper Tuluto Creek. Thickness probably >100 m in some areas.

Dnh NOATAK SANDSTONE AND HUNT FORK SHALE, UNDIVIDED (Upper Devonian)–Transitional facies, consists of brownish-gray micaceous sandstone as beds to 1 m thick as in Noatak Sandstone, interbedded with siltstone and gray silty micaceous shale; forms brownish-gray to yellowish brown weathering slopes in lowlands of western part of map area.

Dhf HUNT FORK SHALE (Upper Devonian)–Dark gray- and olive-green shale, with interbedded thin bedded sandstone, and dominantly micaceous phyllitic shale in lower part. Forms smoothly weathered slopes with a conspicuous micaceous sheen beneath higher ridges of Noatak Sandstone. Thickness >100 m.

Di IGNEOUS ROCKS (Devonian ?)–Basalt or andesite, dark-greenish gray, fine grained, dense; probable sill within Kanayut on south side of Kayakak Mountain in eastern part of map area, and an isolated body within Hunt Fork Shale on north fork of Aniuk River in western part of map, but contacts not exposed.

ANIUK RIVER SEQUENCE, UPPER IPNAVIK PANEL

Dknu KANAYUT CONGLOMERATE AND NOATAK SANDSTONE, UNDIVIDED (Upper Devonian)—Sandstone, minor conglomerate, reddish brown silty mudstone, and phyllitic shale; characterized in places by conspicuous slaty cleavage in sandstone and conglomerate. Generally fine grained quartz sandstone, in units up to 2 m thick, is calcareous, platy, evenly laminated to cross bedded, with conspicuous limonitic spots; interbedded with reddish brown silty mudstone and phyllitic shale. Conglomerate is composed of quartz and black chert pebbles to 1 cm diameter and some red shale rippup clasts, in beds to 2 m thick with prominent trough cross bedding, forms 20 m thick interval. Map unit consists of resistant black-weathering low ridges and tan to reddish-brown weathering slopes separated by tundra, is probably cut by numerous unmapped thrust faults. Forms high mountains between Ipnavik River and Fenian Lake.

Unit probably represents a deeper structural level of the stratigraphically lower part of a transitional marine facies of Kanayut , Noatak , and minor Hunt Fork that has been thrust northward over more proximal upper Kanayut and Noatak, and is apparently structurally overlain by the entire Aniuk River sequence.

CORRELATION OF MAP UNITS

SURFICIAL DEPOSITS

ENDICOTT MOUNTAINS ALLOCHTHON



PICNIC CREEK ALLOCHTHON



IPNAVIK RIVER ALLOCHTHON



COPTER PEAK AND NUKA RIDGE ALLOCHTHONS UNDIVIDED

COPTER PEAK ALLOCHTHON



MISHEGUK MOUNTAIN ALLOCHTHON



MAP SHOWING DISTRIBUTION OF ALLOCHTHONS



STATION AND TRAVERSE MAP



MAP SYMBOLS

- Contact
- High angle fault
- Fault–arrows show relative horizontal movement
- Thrust fault–sawteeth on upper plate dashed where inferred, dotted where concealed
- Major thrust fault bounding allochthons–sawteeth on upper plate, dashed where inferred, dotted where concealed.
- Fold axis showing direction of plunge, dotted where approximate or concealed.
- Anticline
- Syncline
- Strike and dip of beds
  - 70 Inclined bedding
  - Horizontal bedding

REFERENCES CITED

Chapman, R.M., Dotterman, R.L., and Mangus, M.D., 1964, Geologic map and structure sections of the Killik-Etivluk Rivers region, Alaska, in Geology of the Arctic, Proceedings of the First Symposium on Arctic Geology: Arctic Institute of North America, v. 1, p. 448-464, 1 sheet, scale 1:125,000.

Dumoulin, J.A. and Harris, Anita G., 1993, Lithofacies and conodonts of Carboniferous strata in the Ivotuk Hills, western Brooks Range, Alaska, in Bartsch-Bacon, Cynthia, and Till, A.B., eds., Geologic Studies in Alaska by the U.S. Geological Survey, 1992: U.S. Geological Survey Bulletin 2068, p. 41-57.

Dumoulin, J.A., Harrit, Anita G., and Schmidt, J.M., 1994, Deep-water lithofacies and conodonts of Carboniferous strata in the Endicott Mountains allochthon, Alaska in Dumoli-Bacon, Cynthia, and Till, A.B., eds., Geologic Studies in Alaska by the U.S. Geological Survey, 1992: U.S. Geological Survey Bulletin 2068, p. 19-39.

Dutro, J.T., Blodgett, R.B., and Mull, C.G., in press, Cymbospiter from Upper Devonian rocks east of the Endicott Group, west-central Brooks Range, Alaska in 16, A.B., and Moore, T.N., eds., Geologic Studies in Alaska by the U.S. Geological Survey, 1993: U.S. Geological Survey Bulletin 2107, in press.

Ellersieck, Inyo, Tailleur, I.L., and Mull, C. G.,1990, Reconnaissance geologic map of the Story Creek area, National Petroleum Reserve, Alaska: U.S. Geological Survey Open-file Report 90-___, scale 1:63,360.

Hamilton, T.D., 1980, Surficial geologic map of the Killik River quadrangle, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF 1234, 1 sheet, scale 1:250,000.

Hamilton, T.D., 1984, Preliminary surficial geologic map of the Howard Pass quadrangle, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF 1677, 1 sheet, scale 1:250,000.

Mayfield, C.F., Tailleur, I.L., and Ellersieck, Inyo, 1988, Stratigraphy, structure, and paleontological synthesis of the western Brooks Range, northwestern Alaska, in Gryc, George, ed., Geology and Exploration of the National Petroleum Reserve in Alaska, 1974 to 1982: U.S. Geological Survey Professional Paper 1399, p. 143-186.

Moore, T.E., Alexander, J.N., and Werdon, M. (1993) Middle Jurassic U-Pb crystallization age for Siniktanneyuk Mountain ophiolite, Brooks Range, Alaska: Geological Society of America, Cordilleran Section, with Program, v. 5, no.5, p. 124.

Mull, C.G., Crowder, R.K., Adams, K.E., Sisk, J.P., Bodnar, D.A., Harris, E.E., Alexander, R.A., and Solie, D.N., 1987, Stratigraphy and structural setting of the Picnic Creek allochthon, Killik River Quadrangle, central Brooks Range, Alaska, in Tailleur, I.L., and Weimer, Paul, eds., Alaskan North Slope Geology: Pacific Section Society of Economic Paleontologists and Mineralogists, Pacific Section, and Alaska Geological Society, Book 50, p. 659-682.

Mull, C. G., T.E. Moore, E.E. Harris, and I.L. Tailleur, 1994, Geologic Map of the Killik River quadrangle, U.S. Geological Survey Open-file Report, 1 sheet, scale 1:125,000.

Nelson, S.W., and Nelson, W.H. (1982), Geology of the Siniktanneyuk Mountain ophiolite, Howard Pass quadrangle, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF 1441, 1 sheet, scale 1:63,360.

Tailleur, I.L., Kent, B.H., and Nelson, W.H., 1951, Geologic-topologic maps of the Noka Etivluk region, northern Alaska: U.S. Geological Survey Open-file Report OF 90-139, 7 sheets, scale 1:63,360.

SOURCES OF MAP DATA

Unpublished manuscript field maps

Mull, C. G., Moore, T. E., Reilenshuh, R.R., and G.H. Pessel, unpublished geologic mapping in the Killik River AS and part of BS quadrangles, scale 1:63,360.

Mull, C. G., R.R., Reilenshuh, R.R., Moore, T. E. and G.H. Pessel, unpublished geologic mapping in the Killik River AS and part of BS quadrangles, scale 1:63,360.

Mull, C. G., Moore, T.E., and Reilenshuh, R.R., unpublished geologic mapping in parts of the Howard Pass A2 and B2 quadrangles, scale 1:63,360.

Mull, C., E.E. Harris, and Reilenshuh, R.R., unpublished geologic mapping in parts of the Howard Pass A4 and B4 quadrangles, scale 1:63,360.

Mull, C. G., Reilenshuh, R.R, and Dover, J.G., unpublished manuscript geologic mapping in parts of the Howard Pass A4 and B4 quadrangles, scale 1:63,360.

Published maps

Chapman, R.M. and others, 1964.

Ellersieck, Inyo, and others,1990.

Hamilton, T.D., 1980.

Hamilton, T.D., 1984.

Mull, C. G., and others, 1994.

Nelson, S.W., and Nelson, W.H. (1982).

Tailleur, I.L., and others, 1951.


Nimiuk Pass
Mt. Bertie
Funnel Pk.
Ipnavik River
Mountain
Mountain
Mountain