


SURFICIAL GEOLOGIC MAP SHOWING LOCATION OF WELLS AND SPRINGS





MAP SHOWING WATER QUALITY FROM SELECTED SURFACE AND GROUND WATER SOURCES

INTRODUCTION

Development of ground water in the Homer area is limited by objectionable amounts of iron in solution and by small yields available to individual wells. Because surface-water supplies are located in areas that are distant from the population center, development would be costly. In 1960, the U.S. Geological Survey began a reconnaissance study to provide information relating to the water problems of the Homer area. This report briefly discusses the geology of the area and the possibility of future ground-water development. Mapping and descriptions of geologic formations are largely from published reports. Modifications of the formation boundaries in the Homer bench area were made by the senior author. Appreciation is expressed to well owners and drillers for information. F. H. and A. H. Thorn supplied most of the well logs used in interpretation of subsurface geology.

GEOGRAPHIC SETTING

The Homer area lies at the southern end of the Kenai Peninsula between Cook Inlet and the Kenai Mountains. (See index map) The area includes the Homer Spit, which extends 4 miles into Kachemak Bay, two benches, and the adjoining escarpment. The escarpment extends from the mouth of Diamond Creek on the west to Fritz Creek on the east. Altitude of the escarpment ranges from more than 1,000 feet to sea level. North of the escarpment crest the upland lies at altitudes generally higher than 750 feet. The principal features of the area are shown on the oblique aerial photograph. Their relationship to geology is shown on the diagrammatic section.



part. Numerous small creeks that head in the escarpment flow southward into Beluga Mudlake or Kachemak Bay. In the northern part of the area, Bridge Creek and Twitter Creek drain into the Anchor River, the principal drainage system immediately north of the Homer area. The area surrounding Beluga Mudlake and other parts of the lower bench are covered with muskeg and stunted spruce forest. Dense spruce and alder forests occur in scattered areas of the uplands. The population of the area is about 1,200, most of whom are engaged in seasonal commercial

GEOLOGY

The surficial geologic map shows that rocks ranging in age from Tertiary to Recent are exposed within the three principal geomorphic units of the Homer area, the escarpment and adjacent upland, the bench area at the foot of the escarpment, and the Homer Spit. The escarpment is formed on generally northward dipping strata of the Kenai Formation of Tertiary age. The area, at the foot of the escarpment, consists of two benches, a lower bench which extends from about the 50-foot level to the 300-foot level, and an upper bench which extends from the 300-foot to the 400-foot level. The benches are composed of glacial, alluvial-fan, and colluvial deposits of Quaternary age that unconformably overlie Tertiary rocks. The Homer Spit is composed of well-sorted Recent beach deposits overlying marine and glaciolacustrine sediments of Pleistocene age.

TERTIARY SYSTEM

Oligocene(?), Miocene, and Pliocene Series

Kenai Formation

The Kenai Formation, which underlies the entire Homer area, consists of moderately indurated sand, silt, and clay, generally in thin beds and lenses, interbedded with a few thin lenses of fine conglomerate and many thin beds of subbituminous or lignitic coal. Many fine-grained sandstone sequences containing ferruginous masses or ironstone concretions in distinct bands and as scattered nodules are common throughout the formation. The thickness of the Kenai Formation at Homer is unknown. However, an oil-test well drilled by Occidental Petroleum Co., South Diamond Gulch No. 1 (sec. 6, T. 6 S., R. 14 W.), just north west of the study area, drilled through 10, 568 feet of the Kenai Formation. The age of the formation is considered by Wolfe, Hopkins and Leopold (1966) to be late Oligocene(?), Miocene, and Pliocene on the basis of its flora.

QUATERNARY SYSTEM

Pleistocene and Recent Series

The Pleistocene history of the Homer area is quite complex. According to Karlstrom (1964), the area probably has been glaciated several times. The area north of the escarpment crest is covered by deposits of the Elkhina Glaciation; whereas deposits of the Naptovne Glaciation occur in the area east of Homer and along the escarpment face. Most of these deposits are thin and are therefore mapped generally as bedrock of the Kenai Formation after Barnes and Cobb (1959, pl. 17). One of the oldest Quaternary deposits in the area is the Pleistocene moraine deposits of Naptovne Glaciation, which occurs extensively in the eastern part of the study area. The moraines are thin deposits of interbedded sand and gravel. Undifferentiated Quaternary deposits, which are considered to be Pleistocene in age, occur in the northern and western parts of the area. These deposits may contain pre-Naptovne glacial drift that has been eroded and redepos-

ited, probably in Naptovne or later Pleistocene time.

Glaciolacustrine deposits of Pleistocene age range in thickness from less than 10 to more than 60 feet in the bench area near Beluga Mudlake. South of Beluga Mudlake the deposits are represented by coarse well-sorted sand and gravel that appear to have been deposited near ice by running water and later disturbed by ice movement. West of the airport, well records indicate poorly sorted deposits. Where these deposits attain a thickness of more than 20 feet, they have been mapped as subunit 1. Where the deposits are less than 20 feet in thickness, they are mapped as subunit 2. East and north of Beluga Mudlake, the lacustrine clays thin and become indistinguishable from the colluvium and clayey till deposits.



Unexposed deposits of probable Pleistocene age underlie the thin surficial deposits of the Homer Spit. Test drilling on the Spit has indicated about 300 feet of sediments of glaciolacustrine origin. These deposits probably are related, in the upper 200 feet at least, to the thick deposits of glacial material underlying the area at Homer. The lower 100 feet contains increasing amounts of clay with sand and gravel beds.

Two coalescing alluvial fans at the eastern end of Beluga Mudlake consist of sand and gravel of postglacial age. The thickness of this deposit is unknown but probably is small.

Flood-plain deposits which occur in the valley of Fritz Creek and in a few other creeks consist of silt, sand, and gravel of post-glacial age. The deposits are mappable only in the Fritz Creek area; there they are generally less than 5 feet thick.

AVAILABILITY OF WATER

Water in the Homer area, both surface and ground water, is derived from rainfall and snowmelt. Reports by the U.S. Weather Bureau (1961) show a maritime climate characterized by minimum precipitation during the spring months and maximum precipitation during the months of August through December. (See temperature and precipitation graphs). Annual precipitation is about 23 inches at the airport and is about 28 inches on the upland above the escarpment at an altitude of 1,000 feet. During the months of May, June, and July, precipitation is deficient and evaporation and transpiration are believed to be about 5 inches per year, about one-fifth of the total average annual precipitation. No determinations of evaporation and transpiration are available for the Homer area.



Streamflow records are available on the Anchor River 1 mile upstream from the mouth (drainage area 226 square miles). These records show an average discharge for a 12 year period of 298 cfs (cubic feet per second) or 193 mgd (million gallons per day); minimum discharge for the period was 28 cfs or 18 mgd.

The use of surface water at Homer does not appear to be practical at the present time. Use of streamflow would involve long supply lines and possibly reservoirs for the retention of water during winter months. However, additional development in the area may require future consideration of this source of water to supplement ground-water supplies.

Ground Water

The Kenai Formation probably constitutes the most extensive and the most productive aquifer system in the study area. Ground water is available to drilled wells throughout the area. Most of the wells drilled along the bench area yield less than 10 gpm (gallons per minute), although none of the wells drilled were tested to their fullest extent. A larger yield was obtained from well 4 at the Homer hospital, which penetrated 22 feet of sand. The lower 10 feet of the aquifer was screened and the well produced about 20 gpm with about 80 feet of drawdown. This gives a specific capacity of about 0.25 gpm per foot of drawdown, the largest recorded yield per foot of drawdown in the Kenai Formation in the bench area. Most wells drilled in the bench area are not screened and further testing is warranted. Wells drilled into the Kenai Formation in the upland north of the escarpment crest have considerably larger yields. A test well (well 132) adjacent to Bridge Creek, drilled for the city of Homer in 1966, produced about 80 gpm with about

Table of water analyses from selected surface and ground-water sources (Analyses by the U.S. Geological Survey—chemical concentrations in parts per million)

Township range, section, and 1/2 section location	Well no.	Date of collection	Depth of well (feet)	Silica (SiO ₂)	Total iron (Fe)	Manganese (Mn)	Calcium (Ca)	Magnesium (Mg)	Sodium (Na)	Potassium (K)	Bicarbonate (HCO ₃)	Sulfate (SO ₄)	Chloride (Cl)	Fluoride (F)	Dissolved solids (residue on evaporation at 180° C)	Hardness as CaCO ₃	Calc. Magnesium	Non-carbonate	Specific conductance (microhm-cm at 25° C)	pH	Color
6S 18W 18N W/4ESE	4	9-14-62	112	26	0.12 ¹	0.01	8.0	1.0	292	4.4	772	4.0	6.0	0.0	66	258	20	119	8.5	30	
18 N/2ENE	10	12-28-62	195	50	0.09	0.00	6.4	1.0	117	6.3	910	2.0	8.5	2.2	112	48	100	7.0	5	5	
48 4-24-63 Spring	27	02	00	14	2.9	9.6	3.1	72	1.0	8.5	2.2	112	48	100	7.0	5	5	5	5		
19 N/2ENE	5b	6-10-55	107	31	4.4	0.4	0.0	4.8	4.0	2.9	225	7	14	0	4	366	117	300	7.6		
10 N/2EW	11	3-28-62	72	60	6.8	0.0	4.0	4.8	4.0	2.9	225	7	14	0	4	366	117	300	7.6		
14 N/2EW	13	4-24-62	80	21	1.9	0.0	2.4	1.0	40	1.0	400	1.0	16	2	3	744	9	1159	7.2	80	
12 N/2EW	14	12-14-62	107	72	7.1	0.0	4.0	4.8	4.0	2.9	225	7	14	0	4	366	117	300	7.6		
20 N/2EW	30	5-31-53	119	34	2.9	0.0	1.6	2.0	184	2.4	466	0	16	1	1	467	82	201	6.7	0	
15 N/2EW	49a	4-4-60	72	46	11	0.0	2.0	4.2	1.0	1.0	1.0	1.0	1.0	0	154	82	201	6.7	0		
15 N/2EW	49b	4-4-60	247	37	8.5	0.0	4.2	1.0	20	4.3	255	15	12	0	1	383	174	430	6.6	25	
6S 14W 15 N/2W	30	3-25-63	550	48	3.1	0.0	15	1.0	7.0	6.4	112	1.0	6.0	0	0	179	247	7.0	25	0	
6S 13W 1 S/2W	102	4-24-63	115	43	6.8	0.0	4.0	4.8	4.0	2.9	225	7	14	0	4	366	117	300	7.6		
21 S/2W	110	10-25-62	41	33	1.9	1.2	25	11	13	1.7	88	7.0	4.1	0	0	175	108	296	6.2	5	
9 S/2W	6	6-1-66	224	47	10	1.1	10	5.1	5.0	110	1.4	6.4	0.3	144	90	190	7.0	10	10		
Fritz Creek	3-23-62	Creek	30	10 ¹	0.0	11	7.2	7.6	2.5	80	5.0	6.0	0	2	118	23	143	7.1			

¹ In solution at time of analysis.
² Field determination.

80 feet of drawdown, giving a specific capacity of 1 gpm per foot of drawdown. Other tests in this area indicate that withdrawals of 50 to 100 gpm may be possible. Many springs and seeps flow from the exposed beds of the Kenai Formation along the escarpment face above Homer. The average yield of these springs is about 2 gpm. The springs provide additional ground-water recharge to the Quaternary deposits on the lower bench.

The yield of Quaternary deposits of sand and gravel in the glaciolacustrine areas of the lower bench (Q_{gl}) is comparable to the yields of the Kenai Formation. Most wells in these glacial deposits have yields of from 5 to 25 gpm. However, a yield of 70 gpm was reported for well 51 at the Bearfoot Trailer Court in 1959.

QUALITY OF WATER

Ground water in the Homer area is typically of the sodium bicarbonate or calcium bicarbonate type, and in some areas contains as much as 30 ppm (parts per million) iron as well as combustible gas.

Selected chemical analyses, presented on the table of chemical analyses of waters, show that the ground water contains moderate amounts of dissolved solids and is probably suitable for most uses. The hardness ranges from extremely soft (well 30, 4 ppm) to very hard (well 102, 266 ppm). The soft water is generally obtained from the Tertiary Kenai Formation and the harder water from the Quaternary deposits. The pH of the water in the area ranges from 6.7 to 8.5, neither excessively acid nor excessively alkaline. Fluoride is generally found in small amounts but in concentrations too small to have any appreciable effect on the development of tooth enamel in growing children.

PROSPECTS FOR FUTURE DEVELOPMENT

In the past, moderate yields in the Homer area have been obtained from wells in the Kenai Formation and in the Quaternary deposits. Additional ground-water development, particularly of the Kenai Formation on the upland north of the escarpment crest, seems feasible. The city of Homer, in 1966, completed a test-drilling program near Bridge Creek. Yields of 50 to 80 gpm have been developed, and it seems probable that comparable yields can be obtained in the wells in favorable upland areas. The water is of good quality; the water contains only about 1-2

ppm iron (as compared to about 4 ppm from wells in the bench areas) and is low in dissolved solids.

In the future, consideration should be given to the conjunctive use of ground and surface water in the Homer area. The sources could be used alternately depending upon the season of the year. Although the annual range of streamflow is not known, it is probable that low flows during the winter months could be supplemented by the addition of relatively warm ground water which would be advantageous because it would prevent freezing of the distribution lines.

SELECTED REFERENCES

Barnes, F. F., and Cobb, E. H., 1959, Geology and coal resources of the Homer district, Kenai coal field, Alaska; U.S. Geol. Survey Bull. 1058-F, p. 217-260.
Cederstrom, D. J., 1955, Summary of ground-water development in Alaska, 1950; U.S. Geol. Survey Circ. 169, 37 p.
Karlstrom, T. N. V., 1964, Quaternary geology of the Kenai Lowland and glacial history of the Cook Inlet region, Alaska; U.S. Geol. Survey Paper 443, 69 p.
Krisinsky, D. B., 1953, Southwest Kenai Peninsula, Alaska, in Föwé, T. L., and others, Multiple glaciation in Alaska; U.S. Geol. Survey Circ. 289, p. 5-6.
U.S. Public Health Service, 1962, Drinking water standards; Public Health Service Pub. 956, 61 p.
U.S. Weather Bureau, 1961, Climatological data, Alaska annual summary, 1960.
Waller, R. M., 1962, Preliminary report of ground-water conditions at Homer, Alaska; U.S. Geol. Survey open-file rept., 6 p.
1963, Data on wells in the Homer area, Alaska; Alaska Dept. of Health and Welfare Hydrological Data Rept. 23, 24 p.
Wolfe, J. A., Hopkins, D. M., and Leopold, E. B., 1965, Tertiary stratigraphy and paleontology of the Cook Inlet region, Alaska; U.S. Geol. Survey Paper 398-A, 29 p.

WATER RESOURCES AND SURFICIAL GEOLOGY OF THE HOMER AREA, SOUTH-CENTRAL ALASKA

By
Roger M. Waller, Alvin J. Feulner, and
Donald A. Morris