

United States
Department of the Interior
Geological Survey

MULTICHANNEL SEISMIC-REFLECTION PROFILES COLLECTED
IN 1977 IN THE EASTERN GULF OF ALASKA

by

DENNIS M. MANN¹ and TERRY R. BRUNS¹

U.S. Geological Survey
OPEN FILE REPORT
84-6

This report is preliminary and has not been edited or reviewed for conformity with Geological Survey standards and nomenclature.

Any use of trade names and trademarks in this publication is for descriptive purposes only and does not constitute endorsement by the U.S. Geological Survey

1984

¹U.S. Geological Survey
Menlo Park, CA

MULTICHANNEL SEISMIC-REFLECTION PROFILES COLLECTED
IN 1977 IN THE EASTERN GULF OF ALASKA

by
DENNIS M. MANN and TERRY R. BRUNS

During the summer of 1977 the U.S. Geological Survey (USGS) collected approximately 2492 km. of 24-channel seismic-reflection data in the eastern Gulf of Alaska (fig. 1). The profiles were collected on the USGS Research Vessel S.P. Lee using a sound source of five airguns with a volume of 1326 cubic inches of air compressed to approximately 1950 psi. The recording system consisted of a 24-group streamer, 2400 meters long and a GUS (Global Universal Science) model 4200 recording instrument. Shots were fired every 50 meters and the group interval was 100 meters. A 2-millisecond sampling rate was used in the field: the field tapes were later desampled to 4-milliseconds during the demultiplexing process. Navigational control for the survey was by a Marconi integrated satellite-dopplar sonar navigation system. Record lengths vary from 6 to 10 seconds, depending on water depth, to obtain 5 to 6 seconds below the seafloor. Processing was done at the USGS Marine Geology Multichannel Seismic Processing Center in Menlo Park, California, in the sequence: editing-demultiplexing, velocity analysis, stacking, deconvolution-filtering, and finally displayed on an electrostatic plotter. Plate 1 is a trackline chart of the cruise showing location of the line and some shotpoint locations.

- 1) Electrostatically plotted profiles which have been deconvolved and filtered after stacking. Copies of the profiles may be purchased through:

National Geophysical and Solar Terrestrial Data Center
National Oceanic and Atmospheric Administration
Boulder, Colorado 80302

- 2) Digital magnetic stack tapes which have been processed using velocities derived from velocity analysis. These tapes are not deconvolved or frequency filtered. Copies of the stack tapes and a description of the tape format can be obtained at the requesters expense by contacting:

Data Curator
Pacific Branch of Marine Geology
U.S. Geological Survey
345 Middlefield Rd.
Menlo Park, California 94025

- 3) Digital magnetic demultiplexed tapes. These tapes have been edited for missed shots, blanking time, and muting times. Copies of the demultiplexed tapes and a description of the tape formats can be obtained at the requesters expense by contacting the above address.




FIGURE 1. AREA OF STUDY. PLATE I SHOWING DETAILED LOCATION OF TRACKLINES AND SHOTPOINTS